

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
4 October 2001 (04.10.2001)

PCT

(10) International Publication Number  
**WO 01/72728 A2**

(51) International Patent Classification<sup>7</sup>: **C07D 295/185**,  
A61K 31/495, A61P 29/00

Point Road, Groton, CT 06340 (US). **GLADUE, Ronald**,  
Paul [US/US]; Pfizer Global Research & Development,  
Eastern Point Road, Groton, CT 06340 (US).

(21) International Application Number: PCT/IB01/00375

(74) Agents: **LUMB, J., Trevor** et al.; Pfizer Inc., 235 East  
41nd Street, New York, NY 10017 (US).

(22) International Filing Date: 14 March 2001 (14.03.2001)

(25) Filing Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,  
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,  
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,  
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,  
TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(26) Publication Language: English

(30) Priority Data:  
60/193,789 31 March 2000 (31.03.2000) US

(71) Applicant (*for all designated States except US*): **PFIZER  
PRODUCTS INC.** [US/US]; Eastern Point Road, Groton,  
CT 06340 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **BLUMBERG,**  
**Laura, Cook** [US/US]; Pfizer Global Research & De-  
velopment, Eastern Point Road, Groton, CT 06340 (US).  
**BROWN, Matthew, Frank** [US/US]; Pfizer Global Re-  
search & Development, Eastern Point Road, Groton, CT  
06340 (US). **MCGLYNN, Molly, Ann** [US/US]; Pfizer  
Global Research & Development, Eastern Point Road,  
Groton, CT 06340 (US). **POSS, Christopher, Stanley**  
[US/US]; Pfizer Global Research & Development, Eastern

Published:

— *without international search report and to be republished  
upon receipt of that report*


*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*



BEST AVAILABLE COPY

WO 01/72728 A2

(54) Title: NOVEL PIPERAZINE DERIVATIVES



(57) Abstract: A compound of formula (I) or  
the pharmaceutically acceptable salt thereof;  
wherein a, b, c, d, e, j, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, and R<sup>4</sup> are  
as defined above useful to treat inflammation  
and other immune disorders.

## NOVEL PIPERAZINE DERIVATIVES

### Background of the Invention

The present invention relates to novel piperazine derivatives, methods of use and pharmaceutical compositions containing them.


5       The compounds of the invention are potent and selective inhibitors of chemokine binding to its receptor CCR1 found on inflammatory and immunomodulatory cells (preferably leukocytes and lymphocytes). The CCR1 receptor is also sometimes referred to as the CC-CKR1 receptor. These compounds also inhibit MIP-1 $\alpha$  (and the related chemokines shown to interact with CCR1 (e.g., RANTES and MCP-3)) induced chemotaxis of THP-1 cells and  
10 human leukocytes and are potentially useful for the treatment or prevention of autoimmune diseases (such as rheumatoid arthritis, type I diabetes (recent onset), lupus, inflammatory bowel disease, optic neuritis, psoriasis, multiple sclerosis, polymyalgia rheumatica, uveitis, and vasculitis), acute and chronic inflammatory conditions (such as osteoarthritis, adult Respiratory Distress Syndrome, Respiratory Distress Syndrome of infancy, ischemia  
15 reperfusion injury, and glomerulonephritis), allergic conditions (such as asthma and atopic dermatitis), infection associated with inflammation (such as viral inflammation (including influenza and hepatitis) and Guillian-Barre), chronic bronchitis, xeno-transplantation, transplantation tissue rejection (chronic and acute), organ transplant rejection (chronic and acute), atherosclerosis, restenosis, HIV infectivity (co-receptor usage), and granulomatous  
20 diseases (including sarcoidosis, leprosy and tuberculosis) and sequelae associated with certain cancers such as multiple myeloma. Compounds in this series may also limit the production of cytokines at inflammatory sites, including but not limited to TNF and IL-1, as a consequence of decreasing cell infiltration, providing benefit for diseases linked to TNF and IL-1, including congestive heart failure, pulmonary emphysema or dyspnea associated  
25 therewith, emphysema; HIV-1, HIV-2, HIV-3; cytomegalovirus (CMV), adenoviruses, Herpes viruses (*Herpes zoster* and *Herpes simplex*). They may also provide benefit for the sequelae associated with infection where such infection induces production of detrimental inflammatory cytokines such as TNF e.g., fungal meningitis, joint tissue damage, hyperplasia, pannus formation and bone resorption, psoriatic arthritis, hepatic failure, bacterial meningitis,  
30 Kawasaki syndrome, myocardial infarction, acute liver failure, lyme disease, septic shock, cancer, trauma, and malaria.

MIP-1 $\alpha$  and RANTES are soluble chemotactic peptides (chemokines) which are produced by inflammatory cells, in particular CD8<sup>+</sup> lymphocytes, polymorphonuclear leukocytes (PMNs) and macrophages, *J.Biol. Chem.*, **270** (30) 29671-29675 (1995). These  
35 chemokines act by inducing the migration and activation of key inflammatory and immunomodulatory cells. Elevated levels of chemokines have been found in the synovial fluid of rheumatoid arthritis patients, chronic and acute rejecting tissue from transplant patients and

in the nasal secretions of allergic rhinitis patients following allergen exposure (Teran, *et al.*, *J. Immunol.*, 1806-1812 (1996), and Kuna *et al.*, *J. Allergy Clin. Immunol.* 321 (1994)). Antibodies which interfere with the chemokine/receptor interaction by neutralizing MIP1 $\alpha$  or gene disruption have provided direct evidence for the role of MIP-1 $\alpha$  and RANTES in disease by limiting the recruitment of monocytes and CD8+ lymphocytes (Smith *et al.*, *J. Immunol.*, 153, 4704 (1994) and Cook *et al.*, *Science*, 269, 1583 (1995)). Together this data demonstrates that CCR1 receptor antagonists would be an effective treatment of several immune based diseases. The compounds described within are potent and selective antagonists of the CCR1 receptor.

#### Summary of the Invention

The present invention also relates to a compound of the formula



or the pharmaceutically acceptable salt thereof; wherein

a is 1, 2, 3, 4 or 5;

b is 0, 1, 2, 3 or 4;

c is 0 or 1;

d is 1, 2, 3, 4 or 5;

e is 0 or 1;

j is 1, 2, 3, or 4;

X is C(O), C(S) or CH<sub>2</sub>;

Y is CH<sub>2</sub>, or if e is 0, Y is CHR<sup>8</sup> wherein R<sup>8</sup> is hydrogen, (C<sub>6</sub>-C<sub>10</sub>)aryl or NR<sup>9</sup>R<sup>10</sup>;

Z is oxygen, NR<sup>9</sup> or CR<sup>11</sup>R<sup>12</sup>;

each R<sup>1</sup> is independently selected from hydrogen, hydroxy, hydroxysulfonyl, halo, (C<sub>1</sub>-C<sub>6</sub>)alkyl, mercapto, mercapto(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylthio, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylthio(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>6</sub>-C<sub>10</sub>)aryloxy, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, trifluoromethyl, formyl, formyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, nitro, nitroso, cyano, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, trifluoromethoxy, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkylamino, (C<sub>3</sub>-C<sub>7</sub>)cycloalkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>3</sub>-C<sub>7</sub>)cycloalkyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, ((C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyano(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>7</sub>)alkenyl, (C<sub>2</sub>-C<sub>7</sub>)alkynyl, (C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylthio(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino,

- (C<sub>1</sub>-C<sub>6</sub>)alkylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino, (C<sub>6</sub>-C<sub>10</sub>)arylamino, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl((C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonyl, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, carboxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl, (C<sub>6</sub>-C<sub>10</sub>)arylaminocarbonyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, amidino, guanidino, ureido, (C<sub>1</sub>-C<sub>6</sub>)alkylureido, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido, ureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl and (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl;
- each R<sup>2</sup> and R<sup>3</sup> are independently selected from oxo, halo, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>2</sub>-C<sub>6</sub>)alkenyl, H-C(O)-, H-C(O)-(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>2</sub>-C<sub>6</sub>)alkynyl, hydroxy(C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, thio(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyano(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylthio(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylthio(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, azido(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryloxy(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl,

- (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, carboxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, aminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl, (C<sub>6</sub>-C<sub>10</sub>)arylaminocarbonyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, carboxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl/sulfonyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl or R<sup>14</sup>R<sup>15</sup>N(C<sub>1</sub>-C<sub>6</sub>)alkyl wherein R<sup>14</sup> and R<sup>15</sup> are each independently (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl;
- R<sup>4</sup> is (R<sup>5</sup>)<sub>f</sub>(R<sup>6</sup>)<sub>g</sub>(C<sub>6</sub>-C<sub>10</sub>)aryl, (R<sup>5</sup>)<sub>f</sub>(R<sup>6</sup>)<sub>g</sub>(C<sub>3</sub>-C<sub>10</sub>)cycloalkyl, (R<sup>5</sup>)<sub>f</sub>(R<sup>7</sup>)<sub>h</sub>(C<sub>2</sub>-C<sub>9</sub>)heteroaryl, or (R<sup>5</sup>)<sub>f</sub>(R<sup>7</sup>)<sub>h</sub>(C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl, wherein f is 1, 2, 3 or 4; g and h are each independently 0, 1, 2 or 3;
- R<sup>5</sup> is one to three groups independently selected from (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylcarbonyl, (C<sub>2</sub>-C<sub>9</sub>)heteroarylcarbonyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylureido, amino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, acetylamino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, halo(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, acetylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino,

- $C_6$ alkylsulfonylamino, aminocarbonyl( $C_1-C_6$ )alkylsulfonylamino, ( $C_1-C_6$ )alkoxycarbonylamino( $C_1-C_6$ )alkylsulfonylamino, aminosulfonylamino, ( $C_1-C_6$ )alkylaminosulfonylamino, (( $C_1-C_6$ )alkyl)<sub>2</sub>aminosulfonylamino, aminocarbonyl( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkylsulfonylamino, ( $C_2-C_9$ )heterocycloalkyloxycarbonylamino( $C_1-C_6$ )alkylsulfonylamino, ( $C_2-C_9$ )heteroaryloxycarbonylamino( $C_1-C_6$ )alkylsulfonylamino, cyanoguanidino, ( $C_1-C_6$ )alkylcyanoguanidino, (( $C_1-C_6$ )alkyl)<sub>2</sub>cyanoguanidino, ( $C_2-C_9$ )heterocycloalkylcyanoguanidino, ( $C_2-C_9$ )heteroarylcyanoguanidino, ( $C_2-C_9$ )heterocycloalkyl( $C_1-C_6$ )alkylcyanoguanidino, ( $C_2-C_9$ )heteroaryl( $C_1-C_6$ )alkylcyanoguanidino, amino( $C_1-C_6$ )alkylcyanoguanidino, ( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkylcyanoguanidino, (( $C_1-C_6$ )alkyl)<sub>2</sub>amino( $C_1-C_6$ )alkylcyanoguanidino, aminocarbonyl( $C_1-C_6$ )alkylcyanoguanidino, ( $C_1-C_6$ )alkylaminocarbonyl( $C_1-C_6$ )alkylcyanoguanidino, (( $C_1-C_6$ )alkyl)<sub>2</sub>aminocarbonyl( $C_1-C_6$ )alkylcyanoguanidino, aminocarbonyl( $C_1-C_6$ )alkylamino, ( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkylamino, ( $C_1-C_6$ )alkoxycarbonylamino( $C_1-C_6$ )alkylamino, aminosulfonyl( $C_1-C_6$ )alkylamino, ( $C_2-C_9$ )heteroaryl( $C_1-C_6$ )alkylamino, acetylamino( $C_1-C_6$ )alkylamino, (acetyl)(( $C_1-C_6$ )alkyl)amino( $C_1-C_6$ )alkylamino, ( $C_1-C_6$ )alkoxycarbonyl( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkyl, cyano( $C_1-C_6$ )alkylaminoalkyl, aminocarbonyl( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkyl, acetylamino( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkyl, (acetyl)(( $C_1-C_6$ )alkyl)amino( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkoxycarbonylamino( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkyl, ( $C_2-C_9$ )heterocycloalkyloxycarbonylamino( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkyl, ( $C_2-C_9$ )heteroaryloxycarbonylamino( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkyl, cyanoguanidino( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkylcyanoguanidino( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkyl, (( $C_1-C_6$ )alkyl)<sub>2</sub>cyanoguanidino( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkyl, ureido( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkylureido( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkyl, (( $C_1-C_6$ )alkyl)<sub>2</sub>ureido( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkyl, aminocarbonyloxy( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkyl, aminocarbonyl( $C_1-C_6$ )alkylcarbonylamino( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkylaminocarbonyl( $C_1-C_6$ )alkylcarbonylamino( $C_1-C_6$ )alkyl, (( $C_1-C_6$ )alkyl)<sub>2</sub>aminocarbonyl( $C_1-C_6$ )alkylcarbonylamino( $C_1-C_6$ )alkyl, aminosulfonyl( $C_1-C_6$ )alkylcarbonylamino( $C_1-C_6$ )alkyl, ( $C_2-C_9$ )heterocycloalkyloxycarbonylamino( $C_1-C_6$ )alkyl, cyanoguanidino( $C_1-C_6$ )alkylcarbonylamino( $C_1-C_6$ )alkyl, cyano( $C_1-C_6$ )alkylcarbonylamino( $C_1-C_6$ )alkyl, amino( $C_1-C_6$ )alkylaminocarbonylamino( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkylaminocarbonylamino( $C_1-C_6$ )alkyl, (( $C_1-C_6$ )alkyl)<sub>2</sub>amino( $C_1-C_6$ )alkylaminocarbonylamino( $C_1-C_6$ )alkyl, hydroxy( $C_1-C_6$ )alkylaminocarbonylamino( $C_1-C_6$ )alkyl, aminocarbonyl( $C_1-C_6$ )alkylaminocarbonylamino( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkylcarbonylamino( $C_1-C_6$ )alkylaminocarbonylamino( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkylaminocarbonylamino( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkoxycarbonylamino( $C_1-C_6$ )alkylaminocarbonylamino( $C_1-C_6$ )alkyl, ( $C_2-C_9$ )heterocycloalkyloxycarbonylamino( $C_1-C_6$ )alkylaminocarbonylamino( $C_1-C_6$ )alkyl, ( $C_2-C_9$ )heteroaryloxycarbonylamino( $C_1-C_6$ )alkylaminocarbonylamino( $C_1-C_6$ )alkyl,

- $C_8$ )heterocycloalkyloxycarbonylamino( $C_1-C_6$ )alkylaminocarbonylamino( $C_1-C_6$ )alkyl, ( $C_2-C_6$ )heteroaryloxycarbonylamino( $C_1-C_6$ )alkylaminocarbonylamino( $C_1-C_6$ )alkyl, ( $C_2-C_6$ )heterocycloalkyl( $C_1-C_6$ )alkylaminocarbonylamino( $C_1-C_6$ )alkyl, ( $C_2-C_6$ )heteroaryl( $C_1-C_6$ )alkylaminocarbonylamino( $C_1-C_6$ )alkyl, ureido( $C_1-C_6$ )alkylureido( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkylureido( $C_1-C_6$ )alkylureido( $C_1-C_6$ )alkyl, (( $C_1-C_6$ )alkyl)<sub>2</sub>ureido( $C_1-C_6$ )alkylureido( $C_1-C_6$ )alkyl, cyanoguanidino( $C_1-C_6$ )alkylureido( $C_1-C_6$ )alkyl, halo( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkyl, amino( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkyl, (( $C_1-C_6$ )alkyl)<sub>2</sub>amino( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkyl, acetylamino( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkyl, (acetyl)(( $C_1-C_6$ )alkyl)amino( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkyl, ureido( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkylureido( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkyl, (( $C_1-C_6$ )alkyl)<sub>2</sub>ureido( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkyl, cyanoguanidino( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkyl(cyanoguanidino)( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkyl, (( $C_1-C_6$ )alkyl)<sub>2</sub>(cyanoguanidino)( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkyl, aminocarbonyl( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkoxycarbonylamino( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkyl, ( $C_2-C_6$ )heterocycloalkyloxycarbonylamino( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkyl, ( $C_2-C_6$ )heteroaryloxycarbonylamino( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkyl, aminosulfonylamino( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkylaminosulfonylamino( $C_1-C_6$ )alkyl, (( $C_1-C_6$ )alkyl)<sub>2</sub>aminosulfonylamino( $C_1-C_6$ )alkyl, cyanoguanidino( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkyl(cyanoguanidino)( $C_1-C_6$ )alkyl, (( $C_1-C_6$ )alkyl)<sub>2</sub>(cyanoguanidino)( $C_1-C_6$ )alkyl, ( $C_2-C_6$ )heterocycloalkyl(cyanoguanidino)( $C_1-C_6$ )alkyl, ( $C_2-C_6$ )heterocycloalkyl(cyanoguanidino)( $C_1-C_6$ )alkyl, ( $C_2-C_6$ )heteroaryl(cyanoguanidino)( $C_1-C_6$ )alkyl, amino( $C_1-C_6$ )alkyl(cyanoguanidino)( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkyl(cyanoguanidino)( $C_1-C_6$ )alkyl, aminocarbonyl( $C_1-C_6$ )alkyl(cyanoguanidino)( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkylaminocarbonyl( $C_1-C_6$ )alkyl(cyanoguanidino)( $C_1-C_6$ )alkyl, (( $C_1-C_6$ )alkyl)<sub>2</sub>aminocarbonyl( $C_1-C_6$ )alkyl(cyanoguanidino)( $C_1-C_6$ )alkyl, aminosulfonyl, ( $C_1-C_6$ )alkylaminosulfonyl, (( $C_1-C_6$ )alkyl)<sub>2</sub>aminosulfonyl, ( $C_2-C_6$ )heterocycloalkylsulfonyl, amino( $C_1-C_6$ )alkylaminosulfonyl, ( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkylaminosulfonyl, (( $C_1-C_6$ )alkyl)<sub>2</sub>amino( $C_1-C_6$ )alkylaminosulfonyl, ( $C_2-C_6$ )heteroarylaminosulfonyl, hydroxy( $C_1-C_6$ )alkylaminosulfonyl, ( $C_1-C_6$ )alkoxy( $C_1-C_6$ )alkylaminosulfonyl, ureido( $C_1-C_6$ )alkylaminosulfonyl, ( $C_1-C_6$ )alkylureido( $C_1-C_6$ )alkylaminosulfonyl, (( $C_1-C_6$ )alkyl)<sub>2</sub>ureido( $C_1-C_6$ )alkylaminosulfonyl, ( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkylaminosulfonyl, ( $C_1-C_6$ )alkoxycarbonylamino( $C_1-C_6$ )alkylaminosulfonyl, ( $C_2-C_6$ )heterocycloalkyloxycarbonylamino( $C_1-C_6$ )alkylaminosulfonyl, ( $C_2-C_6$ )heteroaryloxycarbonylamino( $C_1-C_6$ )alkylaminosulfonyl, aminocarbonyl( $C_1-C_6$ )alkylaminosulfonyl,

- C<sub>6</sub>alkylaminosulfonyl, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylaminosulfonyl, R<sup>6</sup> is one to three groups independently selected from hydrogen, hydroxy, hydroxysulfonyl, halo, (C<sub>1</sub>-C<sub>6</sub>)alkyl, mercapto, mercapto(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylthio, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl,
- 5 (C<sub>6</sub>-C<sub>10</sub>)arylsulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylthio(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>6</sub>-C<sub>10</sub>)aryloxy, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, trifluoro(C<sub>1</sub>-C<sub>6</sub>)alkyl, formyl, formyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, nitro, nitroso, cyano, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, trifluoro(C<sub>1</sub>-C<sub>6</sub>)alkoxy, amino(C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>3</sub>-C<sub>10</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>10</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>3</sub>-C<sub>10</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>10</sub>)cycloalkylamino,
- 10 (C<sub>3</sub>-C<sub>10</sub>)cycloalkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyano(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (hydroxy) (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkylamino)(C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylthio(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino, (C<sub>6</sub>-C<sub>10</sub>)arylamino, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylamino, (C<sub>2</sub>-C<sub>9</sub>)heteroarylamino, (C<sub>3</sub>-C<sub>10</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino, (C<sub>2</sub>-C<sub>6</sub>)alkenylcarbonylamino, (C<sub>3</sub>-C<sub>10</sub>)cycloalkylcarbonylamino, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonylamino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylcarbonylamino, halo(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, ((C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, C<sub>3</sub>-C<sub>10</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>6</sub>-C<sub>10</sub>)arylsulfonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroarylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl,
- 35 (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl,



- $C_6$ alkylaminocarbonyl,  $((C_1-C_6)alkyl)_2$ aminocarbonyl,  $(C_6-C_{10})$ arylaminocarbonyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkylaminocarbonyl, (aminocarbonyl $(C_1-C_6)$ alkylaminocarbonyl,  $((C_1-C_6)alkylaminocarbonyl(C_1-C_6)alkylaminocarbonyl$ ,  $((C_1-C_6)alkoxycarbonyl(C_1-C_6)alkylaminocarbonyl$ , (amino $(C_1-C_6)alkyl$ )aminocarbonyl, (hydroxy $(C_1-C_6)alkylaminocarbonyl$ , aminocarbonyl $(C_1-C_6)alkyl$ ,  $(C_1-C_6)alkylaminocarbonyl(C_1-C_6)alkyl$ ,  $((C_1-C_6)alkyl)_2$ aminocarbonyl $(C_1-C_6)alkyl$ ,  $(C_6-C_{10})$ arylaminocarbonyl $(C_1-C_6)alkyl$ ,  $(C_6-C_{10})$ aryl $(C_1-C_6)alkylaminocarbonyl(C_1-C_6)alkyl$ , amidino, hydroxyamidino, guanidino, ureido,  $(C_1-C_6)alkylureido$ ,  $(C_6-C_{10})$ arylureido,  $((C_6-C_{10})aryl)_2$ ureido,  $(C_6-C_{10})$ aryl $(C_1-C_6)alkylureido$ , halo $(C_1-C_6)alkylureido$ ,  $((C_1-C_6)alkyl)((C_6-C_{10})aryl)ureido$ ,  $((C_1-C_6)alkyl)_2$ ureido, halo $(C_1-C_6)alkylcarbonylureido$ , ureido $(C_1-C_6)alkyl$ ,  $(C_1-C_6)alkylureido(C_1-C_6)alkyl$ ,  $((C_1-C_6)alkyl)_2$ ureido $(C_1-C_6)alkyl$ ,  $(C_6-C_{10})$ arylureido $(C_1-C_6)alkyl$ ,  $(C_6-C_{10})$ aryl $_2$ ureido $(C_1-C_6)alkyl$ ,  $(C_6-C_{10})$ aryl $(C_1-C_6)alkylureido(C_1-C_6)alkyl$ , halo $(C_1-C_6)alkylureido(C_1-C_6)alkyl$ , (halo $(C_1-C_6)alkyl)((C_1-C_6)alkyl)ureido(C_1-C_6)alkyl$ ,  $((C_1-C_6)alkoxycarbonyl(C_1-C_6)alkyl)ureido(C_1-C_6)alkyl$ , glycinamido,  $(C_1-C_6)alkylglycinamido$ , aminocarbonylglycinamido,  $(C_1-C_6)alkoxy(C_1-C_6)alkylcarbonylglycinamido$ , (aminocarbonyl $((C_1-C_6)alkyl)glycinamido$ ,  $((C_1-C_6)alkoxycarbonyl(C_1-C_6)alkylcarbonyl)((C_1-C_6)alkyl)glycinamido$ ,  $((C_1-C_6)alkoxycarbonylamino(C_1-C_6)alkylcarbonyl)glycinamido$ ,  $(C_6-C_{10})$ arylcarbonylglycinamido,  $((C_6-C_{10})arylcarbonyl)((C_1-C_6)alkyl)glycinamido$ ,  $((C_6-C_{10})aryl(C_1-C_6)alkylaminocarbonyl)glycinamido$ ,  $(C_6-C_{10})$ aryl $(C_1-C_6)alkylaminocarbonyl)((C_1-C_6)alkyl)glycinamido$ ,  $(C_6-C_{10})$ arylaminocarbonylglycinamido,  $((C_6-C_{10})arylaminocarbonyl)((C_1-C_6)alkyl)glycinamido$ , glycinamido $(C_1-C_6)alkyl$ , alaninamido,  $(C_1-C_6)alkylalaninamido$ , alaninamido $(C_1-C_6)alkyl$ ,  $(C_2-C_9)$ heteroaryl,  $(C_2-C_9)$ heterocycloalkyl,  $(C_2-C_9)$ heteroaryl $(C_1-C_6)alkyl$  and  $(C_2-C_9)$ heterocycloalkyl $(C_1-C_6)alkyl$ ;
- $R^7$  is one to three groups independently selected from hydrogen, hydroxy, halo,  $(C_1-C_6)alkyl$ ,  $(C_1-C_6)alkylsulfonyl$ ,  $(C_6-C_{10})$ arylsulfonyl,  $(C_1-C_6)alkoxy$ , hydroxy $(C_1-C_6)alkoxy$ , halo $(C_1-C_6)alkyl$ , fonyl, nitro, cyano, halo $(C_1-C_6)alkoxy$ ,  $(C_2-C_6)alkenyl$ ,  $(C_2-C_6)alkynyl$ ,  $(C_6-C_{10})$ aryl,  $(C_6-C_{10})$ aryl $(C_1-C_6)alkyl$ , amino,  $(C_1-C_6)alkylamino$ ,  $((C_1-C_6)alkyl)_2$ amino,  $(C_6-C_{10})$ arylamino,  $(C_6-C_{10})$ aryl $(C_1-C_6)alkylamino$ ,  $(C_1-C_6)alkylcarbonylamino$ ,  $(C_1-C_6)alkoxycarbonylamino$ ,  $(C_2-C_6)alkenylcarbonylamino$ , cycloalkylcarbonylamino,  $(C_6-C_{10})$ arylcarbonylamino, halo $(C_1-C_6)alkylcarbonylamino$ ,  $(C_1-C_6)alkoxy(C_1-C_6)alkylcarbonylamino$ ,  $(C_1-C_6)alkoxycarbonyl(C_1-C_6)alkylcarbonylamino$ ,  $((C_1-C_6)alkylcarbonyl)((C_1-C_6)alkyl)amino$ ,  $((C_1-C_6)alkoxycarbonyl)((C_1-C_6)alkyl)amino$ ,  $(C_1-C_6)alkylsulfonylamino$ , amino $(C_1-C_6)alkyl$ ,  $(C_1-C_6)alkylamino(C_1-C_6)alkyl$ ,  $((C_1-C_6)alkyl)_2$ amino $(C_1-C_6)alkyl$ ,  $(C_1-C_6)alkylcarbonylamino(C_1-C_6)alkyl$ ,  $(C_6-C_{10})$ arylcarbonylamino $(C_1-C_6)alkyl$ ,  $((C_1-C_6)alkylcarbonyl)((C_1-C_6)alkyl)amino(C_1-C_6)alkyl$ ,  $(C_1-C_6)alkoxycarbonylamino(C_1-C_6)alkyl$ ,  $(C_1-C_6)alkoxycarbonyl$ ,  $(C_6-C_{10})$ aryl $(C_1-C_6)alkoxycarbonyl$ ,  $(C_1-C_6)alkylcarbonyl$ ,  $(C_6-C_{10})$ arylcarbonyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)alkoxycarbonyl$ ,

C<sub>6</sub>alkylcarbonyl, aminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl, (C<sub>6</sub>-C<sub>10</sub>)arylaminocarbonyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, guanidino, ureido, (C<sub>1</sub>-C<sub>6</sub>)alkylureido, ureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, and glycinamido;

5 R<sup>9</sup> and R<sup>10</sup> are each independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl and (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl; and

10 R<sup>11</sup> and R<sup>12</sup> are each independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>3</sub>-C<sub>8</sub>)cycloalkylcarbonylamino, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, ((C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroarylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino, halo(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonylamino, aminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, carboxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl and (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl.

Preferred compounds of formula I include those wherein R<sup>1</sup> is hydrogen, halo, cyano, nitro, trifluoromethyl, trifluoromethoxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy or (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyloxy.

30 Other preferred compounds of formula I include those wherein R<sup>2</sup> and R<sup>3</sup> are each independently selected from (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl.

35 Other preferred compounds of formula I include those wherein c is 1; X is C(O); d is 1; Y is CH<sub>2</sub>; e is 1; and Z is oxygen.

Other preferred compounds of formula I include those wherein c is 1; X is C(O); d is 2; Y is ethylene; and e is 0.

Other preferred compounds of formula I include those wherein c is 1; X is C(O); d is 1; Y is CH<sub>2</sub>; e is 1; and Z is NR<sup>9</sup> wherein R<sup>9</sup> is hydrogen or (C<sub>1</sub>-C<sub>6</sub>)alkyl.

Other preferred compounds of formula I include those wherein c is 1; X is CH<sub>2</sub>; d is 1; Y is CH<sub>2</sub>; e is 1; and Z is oxygen.

5 Other preferred compounds of formula I include those wherein c is 1; X is CH<sub>2</sub>; d is 2; Y is ethylene; and e is 0.

Other preferred compounds of formula I include those wherein c is 1; X is CH<sub>2</sub>; d is 1; Y is CH<sub>2</sub>; e is 1; and Z is NR<sup>9</sup> wherein R<sup>9</sup> is hydrogen or (C<sub>1</sub>-C<sub>6</sub>)alkyl.

10 Other preferred compounds of formula I include those wherein c is 1; X is C(O); d is 1; Y is CHR<sup>8</sup> wherein R<sup>8</sup> is NR<sup>9</sup>R<sup>10</sup>; R<sup>9</sup> and R<sup>10</sup> are each independently hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl; e is 1; and Z is oxygen.

Other preferred compounds of formula I include those wherein c is 1; X is C(O); d is 1; Y is CHR<sup>8</sup> wherein R<sup>8</sup> is NR<sup>9</sup>R<sup>10</sup>; R<sup>9</sup> and R<sup>10</sup> are each independently hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl; e is 1; and Z is CR<sup>11</sup>R<sup>12</sup> wherein R<sup>11</sup> and R<sup>12</sup> are hydrogen.

15 Other preferred compounds of formula I include those wherein c is 1; X is C(O); d is 1; Y is CHR<sup>8</sup> wherein R<sup>8</sup> is NR<sup>9</sup>R<sup>10</sup>; R<sup>9</sup> and R<sup>10</sup> are each independently hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl; e is 1; and Z is NR<sup>9</sup> wherein R<sup>9</sup> is hydrogen or (C<sub>1</sub>-C<sub>6</sub>)alkyl.

Other preferred compounds of formula I include those wherein c is 1; X is CH<sub>2</sub>; d is 1; Y is CHR<sup>8</sup> wherein R<sup>8</sup> is NR<sup>9</sup>R<sup>10</sup>; R<sup>9</sup> and R<sup>10</sup> are each independently hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl; e is 1; and Z is oxygen.

Other preferred compounds of formula I include those wherein c is 1; X is CH<sub>2</sub>; d is 1; Y is CHR<sup>8</sup> wherein R<sup>8</sup> is NR<sup>9</sup>R<sup>10</sup>; R<sup>9</sup> and R<sup>10</sup> are each independently hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl; e is 1; and Z is CR<sup>11</sup>R<sup>12</sup> wherein R<sup>11</sup> and R<sup>12</sup> are hydrogen.

Other preferred compounds of formula I include those wherein c is 1; X is CH<sub>2</sub>; d is 1; Y is CHR<sup>8</sup> wherein R<sup>8</sup> is NR<sup>9</sup>R<sup>10</sup>; R<sup>9</sup> and R<sup>10</sup> are each independently hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl; e is 1; and Z is NR<sup>9</sup> wherein R<sup>9</sup> is hydrogen or (C<sub>1</sub>-C<sub>6</sub>)alkyl.

Other preferred compounds of formula I include those wherein R<sup>4</sup> is (R<sup>5</sup>)<sub>f</sub>(R<sup>6</sup>)<sub>g</sub>(C<sub>6</sub>-C<sub>10</sub>)aryl or (R<sup>5</sup>)<sub>f</sub>(R<sup>7</sup>)<sub>h</sub>(C<sub>2</sub>-C<sub>6</sub>)heteroaryl wherein f, g and h are independently 1 or 2.

Other preferred compounds of formula I include those wherein R<sup>5</sup> is (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylcarbonyl, (C<sub>2</sub>-C<sub>9</sub>)heteroarylcarbonyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl or (C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl.

Other preferred compounds of formula I include those wherein R<sup>5</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino,

(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, or aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino.

- 5 Other preferred compounds of formula I include those wherein R<sup>5</sup> is amino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, acetyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylureido.

- 10 Other preferred compounds of formula I include those wherein R<sup>5</sup> is amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, acetyl(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, aminosulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminosulfonylamino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino or (C<sub>2</sub>-C<sub>9</sub>)heteroaryloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino.

- Other preferred compounds of formula I include those wherein R<sup>5</sup> is cyanoguanidino, (C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>cyanoguanidino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylcyanoguanidino, (C<sub>2</sub>-C<sub>9</sub>)heteroarylcyanoguanidino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, amino(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino or ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino.

- Other preferred compounds of formula I include those wherein R<sup>5</sup> is aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, acetyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino or (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylamino.

Other preferred compounds of formula I include those wherein R<sup>5</sup> is cyano(C<sub>1</sub>-C<sub>6</sub>)alkylaminoalkyl or aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl.

Other preferred compounds of formula I include those wherein R<sup>5</sup> is acetamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)heterocycloalkyloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)heteroaryloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl or aminocarbonyloxy(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl.

Other preferred compounds of formula I include those wherein R<sup>5</sup> is acetamino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)heterocycloalkyloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl or cyano(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl.

Other preferred compounds of formula I include those wherein R<sup>5</sup> is amino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl amino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)heterocycloalkyloxycarbonyl amino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)heteroaryloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl or cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl.

Other preferred compounds of formula I include those wherein R<sup>5</sup> is amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, acetamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkyl,

$C_6$ alkylsulfonylamino( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkyl(cyanoguanidino)( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkyl,  
 $((C_1-C_6)alkyl)_2(cyanoguanidino)(C_1-C_6)alkylsulfonylamino(C_1-C_6)alkyl$ ,  
 aminocarbonyl( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkoxycarbonylamino( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkyl,  
 ( $C_2-C_9$ )heterocycloalkyloxycarbonylamino( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkyl,  
 ( $C_2-C_9$ )heteroaryloxycarbonylamino( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkyl,  
 aminosulfonylamino( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkylaminosulfonylamino( $C_1-C_6$ )alkyl or  $((C_1-C_6)alkyl)_2aminosulfonylamino(C_1-C_6)alkyl$ .

Other preferred compounds of formula I include those wherein  $R^5$  is  
 cyanoguanidino( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkyl(cyanoguanidino)( $C_1-C_6$ )alkyl,  $((C_1-C_6)alkyl)_2(cyanoguanidino)(C_1-C_6)alkyl$ ,  
 ( $C_2-C_9$ )heterocycloalkyl(cyanoguanidino)( $C_1-C_6$ )alkyl, ( $C_2-C_9$ )heteroaryl(cyanoguanidino)( $C_1-C_6$ )alkyl,  
 ( $C_2-C_9$ )heterocycloalkyl( $C_1-C_6$ )alkyl(cyanoguanidino)( $C_1-C_6$ )alkyl, ( $C_2-C_9$ )heteroaryl( $C_1-C_6$ )alkyl(cyanoguanidino)( $C_1-C_6$ )alkyl,  
 amino( $C_1-C_6$ )alkyl(cyanoguanidino)( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkyl(cyanoguanidino)( $C_1-C_6$ )alkyl,  
 $((C_1-C_6)alkyl)_2amino(C_1-C_6)alkyl(cyanoguanidino)(C_1-C_6)alkyl$ ,  
 $C_6$ alkyl, aminocarbonyl( $C_1-C_6$ )alkyl(cyanoguanidino)( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkylaminocarbonyl( $C_1-C_6$ )alkyl(cyanoguanidino)( $C_1-C_6$ )alkyl  
 or  $((C_1-C_6)alkyl)_2aminocarbonyl(C_1-C_6)alkyl(cyanoguanidino)(C_1-C_6)alkyl$ .

Other preferred compounds of formula I include those wherein  $R^5$  is ( $C_2-C_9$ )heterocycloalkylsulfonyl, amino( $C_1-C_6$ )alkylaminosulfonyl, ( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkylaminosulfonyl,  
 $((C_1-C_6)alkyl)_2amino(C_1-C_6)alkylaminosulfonyl$ , ( $C_2-C_9$ )heteroarylaminosulfonyl, ureido( $C_1-C_6$ )alkylaminosulfonyl, ( $C_1-C_6$ )alkylureido( $C_1-C_6$ )alkylaminosulfonyl,  
 $((C_1-C_6)alkyl)_2ureido(C_1-C_6)alkylaminosulfonyl$ , ( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkylaminosulfonyl,  
 ( $C_1-C_6$ )alkoxycarbonylamino( $C_1-C_6$ )alkylaminosulfonyl, ( $C_2-C_9$ )heterocycloalkyloxycarbonylamino( $C_1-C_6$ )alkylaminosulfonyl,  
 ( $C_2-C_9$ )heteroaryloxycarbonylamino( $C_1-C_6$ )alkylaminosulfonyl, aminocarbonyl( $C_1-C_6$ )alkylaminosulfonyl,  
 cyanoguanidino( $C_1-C_6$ )alkylaminosulfonyl, ( $C_2-C_9$ )heteroaryl( $C_1-C_6$ )alkylaminosulfonyl, ( $C_2-C_9$ )heterocycloalkylaminosulfonyl, Other preferred compounds of  
 formula I include those wherein  $R^5$  is halo( $C_1-C_6$ )alkylaminocarbonyl, hydroxy( $C_1-C_6$ )alkylureido, halo( $C_1-C_6$ )alkylsulfonylamino, ( $C_1-C_6$ )alkoxycarbonyl( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkyl,  
 hydroxy( $C_1-C_6$ )alkylaminocarbonylamino( $C_1-C_6$ )alkyl, halo( $C_1-C_6$ )alkylsulfonylamino( $C_1-C_6$ )alkyl, aminosulfonyl, ( $C_1-C_6$ )alkylaminosulfonyl,  
 $((C_1-C_6)alkyl)_2aminosulfonyl$ , hydroxy( $C_1-C_6$ )alkylaminosulfonyl, and ( $C_1-C_6$ )alkoxy( $C_1-C_6$ )alkylaminosulfonyl.

Other preferred compounds of formula I include those wherein  $R^6$  and  $R^7$  are each  
 independently halo, halo( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkoxy, trifluoromethyl, trifluoromethoxy, hydroxy, aminocarbonyl, cyano, ureido, ( $C_1-C_6$ )alkylsulfonylamino, ( $C_1-C_6$ )alkoxycarbonylamino or glycinamino.

The present invention also relates to the pharmaceutically acceptable acid addition salts of compounds of the formula I. The acids which are used to prepare the pharmaceutically acceptable acid addition salts of the aforementioned base compounds of this invention are those which form non-toxic acid addition salts, i.e., salts containing pharmacologically acceptable anions, such as the hydrochloride, hydrobromide, hydroiodide, nitrate, sulfate, bisulfate, phosphate, acid phosphate, acetate, lactate, citrate, acid citrate, tartrate, bitartrate, succinate, maleate, fumarate, gluconate, saccharate, benzoate, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluenesulfonate and pamoate [i.e., 1,1'-methylene-bis-(2-hydroxy-3-naphthoate)]salts.

The invention also relates to base addition salts of formula I. The chemical bases that may be used as reagents to prepare pharmaceutically acceptable base salts of those compounds of formula I that are acidic in nature are those that form non-toxic base salts with such compounds. Such non-toxic base salts include, but are not limited to those derived from such pharmacologically acceptable cations such as alkali metal cations (e.g., potassium and sodium) and alkaline earth metal cations (e.g., calcium and magnesium), ammonium or water-soluble amine addition salts such as N-methylglucamine-(meglumine), and the lower alkanolammonium and other base salts of pharmaceutically acceptable organic amines.

The compounds of this invention may contain olefin-like double bonds. When such bonds are present, the compounds of the invention exist as cis and trans configurations and as mixtures thereof.

The present invention also relates to compounds of formula I wherein any of the hydrogens may optionally be replaced by deuterium.

Unless otherwise indicated, the alkyl, alkenyl and alkynyl groups referred to herein, as well as the alkyl moieties of other groups referred to herein (e.g., alkoxy), may be linear or branched, and they may also be cyclic (e.g., cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl) or be linear or branched and contain cyclic moieties. Unless otherwise indicated, halogen includes fluorine, chlorine, bromine, and iodine.

(C<sub>3</sub>-C<sub>10</sub>)Cycloalkyl when used herein refers to cycloalkyl groups containing zero to two levels of unsaturation such as cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, 1,3-cyclohexadiene, cycloheptyl, cycloheptenyl, bicyclo[3.2.1]octane, norbornyl etc.

(C<sub>2</sub>-C<sub>9</sub>)Heterocycloalkyl when used herein refers to pyrrolidinyl, tetrahydrofuranyl, dihydrofuranyl, tetrahydropyranyl, pyranal, thiopyranal, aziridinyl, oxiranyl, methylenedioxy, chromenyl, barbituryl, isoxazolidinyl, 1,3-oxazolidin-3-yl, isothiazolidinyl, 1,3-thiazolidin-3-yl, 1,2-pyrazolidin-2-yl, 1,3-pyrazolidin-1-yl, piperidinyl, thiomorpholinyl, 1,2-tetrahydrothiazin-2-yl, 1,3-tetrahydrothiazin-3-yl, tetrahydrothiadiazinyl, morpholinyl, 1,2-tetrahydrodiazin-2-yl, 1,3-tetrahydrodiazin-1-yl, tetrahydroazepinyl, piperazinyl, chromanyl, etc.


(C<sub>2</sub>-C<sub>9</sub>)Heteroaryl when used herein refers to furyl, thienyl, thiazolyl, pyrazolyl, isothiazolyl, oxazolyl, isoxazolyl, pyrrolyl, triazolyl, tetrazolyl, imidazolyl, 1,3,5-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,3-oxadiazolyl, 1,3,5-thiadiazolyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, pyridyl, pyrimidyl, pyrazinyl, pyridazinyl, 1,2,4-triazinyl, 1,2,3-triazinyl, 1,3,5-triazinyl, pyrazolo[3,4-b]pyridinyl, cinnolyl, pteridinyl, purinyl, 6,7-dihydro-5H-[1]pyrindinyl, benzo[b]thiophenyl, 5, 6, 7, 8-tetrahydro-quinolin-3-yl, benzoxazolyl, benzothiazolyl, benzisothiazolyl, benzisoxazolyl, benzimidazolyl, thianaphthenyl, isothianaphthenyl, benzofuranyl, isobenzofuranyl, isoindolyl, indolyl, indoliziny, indazolyl, isoquinolyl, quinolyl, phthalazinyl, quinoxaliny, quinazoliny, benzoxazinyl; etc.

10 Aryl when used herein refers to phenyl or naphthyl.


The term "ureido", as used herein, refers to an "amino-carbonyl-amino" moiety.

The term "acetyl", as used herein, refers to an "alkyl-carbonyl" moiety wherein alkyl is defined as above.

15 The term "cyanoguanidino", as used herein, refers to a functional group having the following formula



The term "(C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C=N-CN)amino", as used herein refers to a functional group having the following formula



20 wherein "HET" refers to a (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl or (C<sub>2</sub>-C<sub>9</sub>)heteroaryl group and the nitrogen of said group is the place of attachment.

The term "mercapto", as used herein, refers to a "HS-" moiety.

25 The compounds of this invention include all conformational isomers (e.g., cis and trans isomers) and all optical isomers of compounds of the formula I (e.g., enantiomers and diastereomers), as well as racemic, diastereomeric and other mixtures of such isomers.

The present invention also relates to a pharmaceutical composition for treating or preventing a disorder or condition selected from autoimmune diseases, rheumatoid arthritis, recent onset type I diabetes, lupus, inflammatory bowel disease, optic neuritis, psoriasis, multiple sclerosis, polymyalgia rheumatica, uveitis, vasculitis, acute and chronic inflammatory conditions, osteoarthritis, adult Respiratory Distress Syndrome, Respiratory Distress



Syndrome of infancy, ischemia reperfusion injury, glomerulonephritis, allergic conditions, asthma, atopic dermatitis, infection associated with inflammation, viral inflammation, influenza, hepatitis, Guillian-Barre, chronic bronchitis, xeno-transplantation, chronic and acute transplantation tissue rejection, chronic and acute organ transplant rejection, atherosclerosis, restenosis, HIV infectivity, granulomatous diseases, sarcoidosis, leprosy and tuberculosis and sequelae associated with certain cancers such as multiple myeloma. Compounds in this series may also limit the production of cytokines at inflammatory sites, including but not limited to TNF and IL-1, as a consequence of decreasing cell infiltration, providing benefit for diseases linked to TNF and IL-1, including congestive heart failure, pulmonary emphysema or dyspnea associated therewith, emphysema; HIV-1, HIV-2, HIV-3; cytomegalovirus (CMV), adenoviruses, Herpes viruses (*Herpes zoster* and *Herpes simplex*). They may also provide benefit for the sequelae associated with infection where such infection induces production of detrimental inflammatory cytokines such as TNF e.g, fungal meningitis, joint tissue damage, hyperplasia, pannus formation and bone resorption, psoriatic arthritis, hepatic failure, bacterial meningitis, Kawasaki syndrome, myocardial infarction, acute liver failure, lyme disease, septic shock, cancer, trauma, and malaria in a mammal, preferably a human, comprising an amount of a compound of the formula I or a pharmaceutically acceptable salt thereof effective in treating or preventing such disorder or condition and a pharmaceutically acceptable carrier.

The present invention also relates to a pharmaceutical composition for treating or preventing a disorder or condition that can be treated or prevented by inhibiting chemokine binding to the receptor CCR1 in a mammal, preferably a human, comprising an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, effective in treating or preventing such disorder or condition and a pharmaceutically acceptable carrier. Examples of such disorders and conditions are those enumerated in the preceding paragraph.

The present invention also relates to a method for treating or preventing a disorder or condition selected from autoimmune diseases, rheumatoid arthritis, recent onset type I diabetes, lupus, inflammatory bowel disease, optic neuritis, psoriasis, multiple sclerosis, polymyalgia rheumatica, uveitis, vasculitis, acute and chronic inflammatory conditions, osteoarthritis, adult Respiratory Distress Syndrome, Respiratory Distress Syndrome of infancy, ischemia reperfusion injury, glomerulonephritis, allergic conditions, asthma, atopic dermatitis, infection associated with inflammation, viral inflammation, influenza, hepatitis, Guillian-Barre, chronic bronchitis, xeno-transplantation, chronic and acute transplantation tissue rejection, chronic and acute organ transplant rejection, atherosclerosis, restenosis, HIV infectivity, granulomatous diseases, sarcoidosis, leprosy and tuberculosis and sequelae associated with certain cancers such as multiple myeloma. Compounds in this series may also limit the production of cytokines at inflammatory sites, including but not limited to TNF and IL-1, as a consequence of decreasing cell infiltration, providing benefit for diseases linked

to TNF and IL-1, including congestive heart failure, pulmonary emphysema or dyspnea associated therewith, emphysema; HIV-1, HIV-2, HIV-3; cytomegalovirus (CMV), adenoviruses, Herpes viruses (*Herpes zoster* and *Herpes simplex*). They may also provide benefit for the sequelae associated with infection where such infection induces production of detrimental inflammatory cytokines such as TNF e.g, fungal meningitis, joint tissue damage, hyperplasia, pannus formation and bone resorption, psoriatic arthritis, hepatic failure, bacterial meningitis, Kawasaki syndrome, myocardial infarction, acute liver failure, lyme disease, septic shock, cancer, trauma, and malaria in a mammal, preferably a human, comprising administering to a mammal in need of such treatment or prevention an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder or condition.

The present invention also relates to a method for treating or preventing a disorder or condition that can be treated or prevented by antagonizing the CCR1 receptor in a mammal, preferably a human, comprising administering to a mammal in need of such treatment or prevention an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder or condition.

The present invention also relates to a pharmaceutical composition for treating or preventing a disorder or condition selected from autoimmune diseases, rheumatoid arthritis, recent onset type I diabetes, lupus, inflammatory bowel disease, optic neuritis, psoriasis, multiple sclerosis, polymyalgia rheumatica, uveitis, vasculitis, acute and chronic inflammatory conditions, osteoarthritis, adult Respiratory Distress Syndrome, Respiratory Distress Syndrome of Infancy, ischemia reperfusion injury, glomerulonephritis, allergic conditions, asthma, atopic dermatitis, infection associated with inflammation, viral inflammation, influenza, hepatitis, Guillian-Barre, chronic bronchitis, xeno-transplantation, chronic and acute transplantation tissue rejection, chronic and acute organ transplant rejection, atherosclerosis, restenosis, HIV infectivity, granulomatous diseases, sarcoidosis, leprosy and tuberculosis and sequelae associated with certain cancers such as multiple myeloma. Compounds in this series may also limit the production of cytokines at inflammatory sites, including but not limited to TNF and IL-1, as a consequence of decreasing cell infiltration, providing benefit for diseases linked to TNF and IL-1, including congestive heart failure, pulmonary emphysema or dyspnea associated therewith, emphysema; HIV-1, HIV-2, HIV-3; cytomegalovirus (CMV), adenoviruses, Herpes viruses (*Herpes zoster* and *Herpes simplex*). They may also provide benefit for the sequelae associated with infection where such infection induces production of detrimental inflammatory cytokines such as TNF e.g, fungal meningitis, joint tissue damage, hyperplasia, pannus formation and bone resorption, psoriatic arthritis, hepatic failure, bacterial meningitis, Kawasaki syndrome, myocardial infarction, acute liver failure, lyme disease, septic shock, cancer, trauma, and malaria in a mammal, preferably a human, comprising a CCR1

receptor antagonizing effective amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

The present invention also relates to a pharmaceutical composition for treating or preventing a disorder or condition that can be treated or prevented by antagonizing the CCR1  
5 receptor in a mammal, preferably a human, comprising a CCR1 receptor antagonizing effective amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

The present invention also relates to a method for treating or preventing a disorder or condition selected from autoimmune diseases, rheumatoid arthritis, recent onset type I  
10 diabetes, lupus, inflammatory bowel disease, optic neuritis, psoriasis, multiple sclerosis, polymyalgia rheumatica, uveitis, vasculitis, acute and chronic inflammatory conditions, osteoarthritis, adult Respiratory Distress Syndrome, Respiratory Distress Syndrome of infancy, ischemia reperfusion injury, glomerulonephritis, allergic conditions, asthma, atopic dermatitis, infection associated with inflammation, viral inflammation, influenza, hepatitis,  
15 Guillian-Barre, chronic bronchitis, xeno-transplantation, chronic and acute transplantation tissue rejection, chronic and acute organ transplant rejection, atherosclerosis, restenosis, HIV infectivity, granulomatous diseases, sarcoidosis, leprosy and tuberculosis and sequelae associated with certain cancers such as multiple myeloma. Compounds in this series may also limit the production of cytokines at inflammatory sites, including but not limited to TNF  
20 and IL-1, as a consequence of decreasing cell infiltration, providing benefit for diseases linked to TNF and IL-1, including congestive heart failure, pulmonary emphysema or dyspnea associated therewith, emphysema; HIV-1, HIV-2, HIV-3; cytomegalovirus (CMV), adenoviruses, Herpes viruses (*Herpes zoster* and *Herpes simplex*). They may also provide benefit for the sequelae associated with infection where such infection induces production of  
25 detrimental inflammatory cytokines such as TNF e.g. fungal meningitis, joint tissue damage, hyperplasia, pannus formation and bone resorption, psoriatic arthritis, hepatic failure, bacterial meningitis, Kawasaki syndrome, myocardial infarction, acute liver failure, Lyme disease, septic








- C<sub>6</sub>)alkylcarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, acetylamino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, aminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (amino(C<sub>1</sub>-C<sub>6</sub>)alkyl)ureido, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, acetylamino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, carboxy(C<sub>1</sub>-C<sub>6</sub>)alkylureido, halo(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, acetylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, aminosulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminosulfonylamino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>2</sub>-C<sub>9</sub>)heteroaryloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, cyanoguanidino, (C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>cyanoguanidino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylcyanoguanidino, (C<sub>2</sub>-C<sub>9</sub>)heteroarylcyanoguanidino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, amino(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylamino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, carboxy(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, acetylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino, (C<sub>6</sub>-C<sub>10</sub>)arylamino

- (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>2</sub>-C<sub>6</sub>)heterocycloalkylamino, (C<sub>2</sub>-C<sub>6</sub>)heteroarylamino, (C<sub>3</sub>-C<sub>10</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino, (C<sub>2</sub>-C<sub>6</sub>)alkenylcarbonylamino, (C<sub>3</sub>-C<sub>10</sub>)cycloalkylcarbonylamino, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonylamino, (C<sub>2</sub>-C<sub>6</sub>)heterocycloalkylcarbonylamino, halo(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, ((C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>3</sub>-C<sub>10</sub>)cycloalkylamino, ureido, (C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>6</sub>-C<sub>10</sub>)arylureido, ((C<sub>6</sub>-C<sub>10</sub>)aryl)<sub>2</sub>ureido, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, halo(C<sub>1</sub>-C<sub>6</sub>)alkylureido, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)((C<sub>6</sub>-C<sub>10</sub>)aryl)ureido, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido, halo(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylureido, glycinamido, (C<sub>1</sub>-C<sub>6</sub>)alkylglycinamido, aminocarbonylglycinamido, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylglycinamido, (aminocarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)glycinamido, ((C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)glycinamido, ((C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)glycinamido, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonylglycinamido, ((C<sub>6</sub>-C<sub>10</sub>)arylcabonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)glycinamido, ((C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl)glycinamido, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)glycinamido, (C<sub>6</sub>-C<sub>10</sub>)arylaminoacarbonylglycinamido and ((C<sub>6</sub>-C<sub>10</sub>)arylaminoacarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)glycinamido.


- R<sup>18</sup> and R<sup>19</sup> together with the nitrogen to which they are attached is selected from the group consisting of (C<sub>2</sub>-C<sub>6</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>2</sub>-C<sub>6</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylamino, halo(C<sub>1</sub>-C<sub>6</sub>)alkylamino, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, carboxy(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, cyano(C<sub>1</sub>-C<sub>6</sub>)alkylamino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, acetyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>2</sub>-C<sub>6</sub>)heterocycloalkyloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>2</sub>-C<sub>6</sub>)heteroaryloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylamino, aminocarbonyloxy(C<sub>1</sub>-C<sub>6</sub>)alkylamino, acetyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>2</sub>-

- C<sub>9</sub>)heterocycloalkyloxycarbonylamino, (C<sub>2</sub>-C<sub>9</sub>)heteroarylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, cyano(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino-carbonylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl amino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino, carboxy(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl amino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino, (C<sub>1</sub>-C<sub>6</sub>) alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl amino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyloxycarbonyl amino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino, (C<sub>2</sub>-C<sub>9</sub>)heteroarylloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylureido, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylureido, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, acetylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>2</sub>-C<sub>9</sub>)heteroarylloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, aminosulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminosulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyanoguanidino, (C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino), ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>(cyanoguanidino), (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(cyanoguanidino), (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(cyanoguanidino), (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(cyanoguanidino), (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino), (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino), amino(C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino), (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino), ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino), aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino), (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino), ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino), (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>2</sub>-C<sub>9</sub>)heteroarylamino, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>2</sub>-C<sub>9</sub>)heteroarylloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino,



- C<sub>8</sub>)alkylamino, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, halo(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino, ureido, (C<sub>1</sub>-C<sub>6</sub>)alkylureido, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido, amino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>3</sub>-C<sub>10</sub>)cycloalkylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>6</sub>-C<sub>10</sub>)arylamino, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, (C<sub>3</sub>-C<sub>10</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, (C<sub>6</sub>-C<sub>10</sub>)arylsulfonylamino, ((C<sub>6</sub>-C<sub>10</sub>)arylsulfonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylamino, (C<sub>2</sub>-C<sub>9</sub>)heteroarylamino, halo(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>6</sub>-C<sub>10</sub>)arylamino, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (carboxy(C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, ((C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (amino(C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, (hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylamino, ureido, (C<sub>1</sub>-C<sub>6</sub>)alkylureido, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido, (C<sub>6</sub>-C<sub>10</sub>)arylureido, (C<sub>6</sub>-C<sub>10</sub>)aryl)<sub>2</sub>ureido, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, halo(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (halo(C<sub>1</sub>-C<sub>6</sub>)alkyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)ureido, ((C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl)ureido, and glycinamido.

PREPARATION A





PREPARATION B


XXII




XXI





XX




XIX

PREPARATION B (Con't)


XIX





XVIII

PREPARATION C



SCHEME 1




XXX




X




I

SCHEME 2

## SCHEME 3




## SCHEME 4




-31-

## SCHEME 5









## SCHEME 6




## SCHEME 7



In reaction 1 of Preparation A, the compound of formula **XXXV**, wherein b is 0, 1 or 2, is converted to the corresponding compound of formula **XXXIV** by reacting **XXXV** with an ethyldiamine compound of the formula,  $(R^3)_j$ -ethyldiamine, in the presence of an aprotic solvent, such as diethylether. The reaction mixture is heated to reflux for a time period  
 5 between about 1 hour to about 12 hours.


In reaction 2 of Preparation A, the compound of formula **XXXIV** is converted to the corresponding compound of formula **XXXIII** by reducing **XXXIV** with a reducing agent, such as sodium borohydride, in a refluxing protic solvent, such as ethanol.

In reaction 3 of Preparation A, the compound of formula **XXXIII** is converted to the  
 10 corresponding compound of formula **XXX** by reacting **XXXIII** with a benzaldehyde compound of the formula




in the presence of a base, such as triethylamine, and a reducing agent, such as sodium triacetoxyborohydride, in an aprotic solvent, such as 1,2-dichloroethane. The reaction mixture  
 15 is stirred at room temperature for a time period between about 1 hour to about 4 hours, preferably about 2 hours.

In reaction 1 of Preparation B, the compound of formula **XXII**, wherein b is 0, 1 or 2, is converted to the corresponding compound of formula **XXI** by reacting **XXIII** with a benzaldehyde compound of the formula



20 in the presence of a base, such as triethylamine, a reducing agent, such as sodium borohydride and an aprotic solvent, such as 1,2-dichloroethane. The reaction is stirred, at room temperature, for a time period between about 1 hour to about 4 hours, preferably about 2 hours.

25 In reaction 2 of Preparation B, the compound of formula **XXI** is converted to the corresponding compound of formula **XX** by first reacting a compound of the formula



wherein *j* is 0, 1 or 2, with 4-methyl morpholine and isobutylchloroformate in the presence of a polar aprotic solvent, such as tetrahydrofuran, followed by reacting the intermediate so formed with the compound of formula **XXI**. The reaction mixture, so formed, is stirred overnight at room temperature.

5 In reaction 3 of Preparation B, the compound of formula **XX** is converted to the corresponding piperazine-2,5-dione compound of formula **XIX** by treating **XX** with trifluoroacetic acid in the presence of a polar aprotic solvent, such as methylene chloride. The reaction is stirred, at room temperature, for a time period between about 1 hour to about 4 hours, preferably about 2 hours.

10 In reaction 4 of Preparation B, the compound of formula **XIX** is converted to the corresponding compound of formula **XVIII** by reducing **XIX** with a reducing agent, such as lithium aluminum hydride. The reaction is conducted at a temperature between about -10°C to about 10°C, preferably about 0°C, for a time period between about 10 minutes to about 90 minutes, preferably about 40 minutes.

15 In reaction 1 of the Preparation C, the compound of formula **XXV** is converted to the corresponding compound of formula **XXIV** by reacting **XXV** with an amine of the formula,  $\text{NHR}^{18}\text{R}^{19}$ , wherein  $\text{R}^{18}$  and  $\text{R}^{19}$  are each independently selected from hydrogen, a nitrogen containing  $(\text{C}_2\text{-C}_9)$ heterocycloalkyl or  $(\text{C}_2\text{-C}_9)$ heteroaryl group, or  $(\text{C}_1\text{-C}_6)$ alkyl optionally substituted by hydroxy, aminocarbonyl,  $(\text{C}_1\text{-C}_6)$ alkylaminocarbonyl,  $((\text{C}_1\text{-C}_6)\text{alkyl})_2\text{carbonyl}$ ,  
20 carboxy,  $(\text{C}_1\text{-C}_6)$ alkylsulfonylamino,  $(\text{C}_1\text{-C}_6)$ alkoxycarbonylamino, aminosulfonyl,  $(\text{C}_1\text{-C}_6)$ alkylaminosulfonyl,  $((\text{C}_1\text{-C}_6)\text{alkyl})_2\text{aminosulfonyl}$ ,  $(\text{C}_8\text{-C}_{10})$ alkoxy,  $(\text{C}_2\text{-C}_9)$ heteroaryl,  $(\text{C}_2\text{-C}_9)$ heterocycloalkyl,  $(\text{C}_1\text{-C}_6)$ alkylcarbonylamino,  $((\text{C}_1\text{-C}_6)\text{alkylcarbonyl})((\text{C}_1\text{-C}_6)\text{alkyl})\text{amino}$ , cyano, ureido,  $(\text{C}_1\text{-C}_6)$ alkylureido,  $((\text{C}_1\text{-C}_6)\text{alkyl})_2\text{ureido}$ , cyanoguanidino,  $(\text{C}_1\text{-C}_6)$ alkylcyanoguanidino and  $((\text{C}_1\text{-C}_6)\text{alkyl})_2\text{cyanoguanidino}$ , or  $\text{R}^{18}$  and  $\text{R}^{19}$  are taken together  
25 with the nitrogen to which they are attached to form a  $(\text{C}_2\text{-C}_9)$ heteroaryl or  $(\text{C}_2\text{-C}_9)$ heterocycloalkyl group, in the presence of a polar aprotic solvent, such as methylene chloride. The reaction mixture is stirred, at room temperature, for a time period between about 1 hour to about 24 hours, preferably about 12 hours.

In reaction 2 of Preparation C, the compound of formula **XXIV** is converted to the  
30 corresponding compound of formula **XXIII** by reacting **XXIV** with thiophenol in the presence of a base, such as sodium hydride, and a polar aprotic solvent, such as dimethylformamide. The reaction is heated to reflux for a time period between about 1 hour to about 10 hours, preferably about 4 hours.

In reaction 3 of Preparation C, the compound of formula **XXV** is converted to the  
35 corresponding compound of formula **XXXVIII** by reacting **XXV** with sodium cyanate in the presence of pyridine and a polar aprotic solvent, such as acetonitrile. The reaction is stirred, at room temperature, for a time period between about 2 hours to about 18 hours, preferably

about 10 hours. An amine of the formula,  $H_2N-C(O)-NR^{18}R^{19}$ , is then added and the reaction mixture so formed is stirred, at room temperature, for a time period between about 2 hours to about 24 hours, preferably about 8 hours.

In reaction 4 of Preparation C, the compound of formula XXXVIII is converted to the  
5 corresponding compound of formula XXXVII according to the procedure described above in reaction 2 of Preparation C.

In reaction 1 of Scheme 1, the compound of formula XXX is converted to the  
corresponding compound of formula X by reacting XXX with a compound of the formula,  $A-(X)_c-(Y)_d-A$ , wherein A is chloro or bromo, in the presence of a base, such as triethylamine,  
10 and a polar aprotic solvent, such as methylene chloride. The reaction is stirred at a temperature between about  $-10^\circ\text{C}$  to about  $10^\circ\text{C}$ , for a time period between about 15 minutes to about 90 minutes, preferably about 30 minutes.

In reaction 2 of Scheme 1, the compound of formula X is converted to the  
corresponding compound of formula I by reacting X with a compound of the formula,  $H-(Z)_e-R^4$  wherein e is 1 and Z is oxygen, in the presence of potassium carbonate, potassium iodide  
15 and an aprotic solvent, such as butanone. The reaction is heated to reflux for a time period between about 4 hours to about 8 hours, preferably about 6 hours.

In reaction 1 of Scheme 2, the compound of formula XXX is converted to the  
corresponding compound of formula I by reacting XXX with a compound of the formula,  $A-(X)_c-(Y)_d-(Z)_e-R^4$ , wherein A is chloro or bromo, in the presence of a base, such as  
20 triethylamine, and a polar aprotic solvent, such as methylene chloride. The reaction is stirred at a temperature between about  $-10^\circ\text{C}$  to about  $10^\circ\text{C}$ , for a time period between about 15 minutes to about 90 minutes, preferably about 30 minutes.

In reaction 1 of Scheme 3, the compound of formula X is converted to the  
25 corresponding compound of formula XII according to the procedure described above in reaction 2 of Scheme 1.

In reaction 2 of Scheme 3, the compound of formula XII is converted to the  
corresponding compound of formula XI by reacting XII with lithium hydroxide monohydrate in  
the presence of methanol, tetrahydrofuran and water. The reaction mixture is stirred  
30 overnight at room temperature.

In reaction 3 of Scheme 3, the compound of formula XI is converted to the  
corresponding compound of formula II, by reacting XI with an amine, in the presence of 4-  
dimethylaminopyridine, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide and a polar aprotic  
solvent, such as methylene chloride. The resulting reaction mixture is stirred overnight at  
35 room temperature.

In reaction 1 of Scheme 4, the compound of formula **X** is converted to the corresponding compound of formula **XV** according to the procedure described above in reaction 2 of Scheme 1.

5 In reaction 2 of Scheme 4, the compound of formula **XV** is converted to the corresponding compound of formula **XIV** by hydrogenating **XV** in the presence of a catalyst, such as platinum on carbon, and a polar protic solvent, such as ethanol. The reaction is carried out under a pressure between about 30 psi to about 40 psi, preferably about 35 psi, for a time period between about 15 minutes to about 1 hour, preferably 30 minutes.

10 In reaction 3 of Scheme 4, for urea formation, the compound of formula **XIV** is converted to the corresponding compound of formula **V** by first reacting **XIV** with 4-nitrophenyl chloroformate in the presence of a base, such as pyridine, and a polar aprotic solvent, such as methylene chloride, followed by reacting the intermediate so formed with an amine. The reaction mixture, so formed, is allowed to stir overnight at room temperature. For sulfonamide formation, the compound of formula **XIV** is reacted with a sulfonyl chloride compound of the  
15 formula,  $R^{18}-Cl$ , in the presence of a base, such as triethylamine, and a polar aprotic solvent, such as methylene chloride. The reaction is stirred overnight at ambient temperature. For cyanoguanidine formation, the compound of formula **XIV** is first treated with sodium hydride in an aprotic solvent, such as tetrahydrofuran, followed by reacting, the intermediate so formed with dimethyl-N-cyanodithio iminocarbonate. The reaction mixture so formed is heated to  
20 reflux overnight. The N-cyano-S-methyl-isothiurea intermediate is then reacted with an amine in the presence of a polar protic solvent, such as methanol. For amide formation, the compound of formula **XIV** is reacted with an acid, such as 3-tert-butoxycarbonylaminopropionic acid in the presence of N-methylmorpholine, O-benzotriazole-1-yl-N,N,N', N'-tetramethyluronium hexafluorophosphate and a polar aprotic solvent, such as  
25 methylene chloride.

In reaction 1 of Scheme 5, the compound of formula **X** is converted to the corresponding compound of formula **XVI**, wherein k is 0, 1, 2, 3 or 4, according to the procedure described above in reaction 2 of Scheme 1.

30 In reaction 2 of Scheme 5, the compound of formula **XVI** is converted to the corresponding compound of formula **VII** by reacting **XVI** with an amine of the formula,  $R^{16} R^{17}N$ , wherein  $R^{16}$  and  $R^{17}$  are each independently hydrogen, a nitrogen containing  $(C_2-C_9)$ heterocycloalkyl or  $(C_2-C_9)$ heteroaryl group, or  $(C_1-C_6)$ alkyl optionally substituted by hydroxy, aminocarbonyl,  $(C_1-C_6)$ alkylaminocarbonyl,  $((C_1-C_6)alkyl)_2$ carbonyl, carboxy,  $(C_1-C_6)alkylsulfonylamino$ ,  $(C_1-C_6)alkoxycarbonylamino$ , aminosulfonyl,  $(C_1-C_6)alkylaminosulfonyl$ ,  
35  $((C_1-C_6)alkyl)_2$ aminosulfonyl,  $(C_6-C_{10})alkoxy$ ,  $(C_2-C_9)heteroaryl$ ,  $(C_2-C_9)heterocycloalkyl$ ,  $(C_1-C_6)alkylcarbonylamino$ ,  $((C_1-C_6)alkylcarbonyl)((C_1-C_6)alkyl)amino$ , cyano, ureido,  $(C_1-C_6)alkylureido$ ,  $((C_1-C_6)alkyl)_2$ ureido, cyanoguanidino,  $(C_1-C_6)alkylcyanoguanidino$  and  $((C_1-$

$C_6$ alkyl)<sub>2</sub>cyanoguanidino, in the presence of a 10:1 ratio solution of dichloroethane/acetic acid. The reaction mixture is stirred, at room temperature, for a time period between about 30 minutes to about 2 hours, preferably about 1 hour. A reducing agent, such as sodium cyanoborohydride is then added to the mixture and the reaction is allowed to stir overnight at room temperature. When R<sup>16</sup> and/or R<sup>17</sup> is/are hydrogen, the compound of formula VII may further be reacted according to the procedure described above in reaction 3 of Scheme 4, to provide ureas, sulfonamides, cyanoguanidinos, or amides.

In reaction 1 of Scheme 6, the compound of formula X is converted to the corresponding compound of formula XXXIX according to the procedure described above in reaction 2 of Scheme 1.

In reaction 2 of Scheme 6, the compound of formula X is converted to the corresponding compound of formula XXXX according to the procedure described above in reaction 2 of Scheme 1.

In reaction 1 of Scheme 7, the acid compound of formula XXXVI is converted to the corresponding compound of formula XXXII by treating XXXVI with thionyl chloride neat or in an aprotic solvent, at room temperature, for a time period between about 1 hour to about 24 hours, preferably 1 hour. The acid chloride so formed is dissolved in a polar aprotic solvent with a compound of the formula, (H<sub>3</sub>CO)(H<sub>3</sub>C)NH•HCl, in the presence of an amine base, such as triethylamine. The reaction mixture is stirred, at room temperature, for a time period between about 1 hour to about 48 hours, preferably about 12 hours.

In reaction 2 of Scheme 7, the amide compound of formula XXXII is converted to the corresponding compound of formula XXXI by reacting XXXII with a (C<sub>2</sub>-C<sub>9</sub>)heteroaryl lithium reagent in the presence of a polar aprotic solvent at a temperature between about -100°C to room temperature, preferably about -78°C. The resulting reaction mixture is stirred for a time period between about 1 hour to about 24 hours, preferably about 12 hours, at a temperature between about -78°C to about 50°C, preferably about 20°C.

Unless indicated otherwise, the pressure of each of the above reactions is not critical. Generally, the reactions will be conducted at a pressure of about one to about three atmospheres, preferably at ambient pressure (about one atmosphere).

The compounds of the formula I which are basic in nature are capable of forming a wide variety of different salts with various inorganic and organic acids. Although such salts must be pharmaceutically acceptable for administration to animals, it is often desirable in practice to initially isolate a compound of the formula I from the reaction mixture as a pharmaceutically unacceptable salt and then simply convert the latter back to the free base compound by treatment with an alkaline reagent, and subsequently convert the free base to a pharmaceutically acceptable acid addition salt. The acid addition salts of the base compounds of this invention are readily prepared by treating the base compound with a

substantially equivalent amount of the chosen mineral or organic acid in an aqueous solvent medium or in a suitable organic solvent such as methanol or ethanol. Upon careful evaporation of the solvent, the desired solid salt is obtained.

The acids which are used to prepare the pharmaceutically acceptable acid addition salts of the base compounds of this invention are those which form non-toxic acid addition salts, i.e., salts containing pharmacologically acceptable anions, such as hydrochloride, hydrobromide, hydroiodide, nitrate, sulfate or bisulfate, phosphate or acid phosphate, acetate, lactate, citrate or acid citrate, tartrate or bitartrate, succinate, maleate, fumarate, gluconate, saccharate, benzoate, methanesulfonate and pamoate [i.e., 1,1'-methylene-bis-(2-hydroxy-3-naphthoate)] salts.

Those compounds of the formula I which are also acidic in nature, are capable of forming base salts with various pharmacologically acceptable cations. Examples of such salts include the alkali metal or alkaline-earth metal salts and particularly, the sodium and potassium salts. These salts are all prepared by conventional techniques. The chemical bases which are used as reagents to prepare the pharmaceutically acceptable base salts of this invention are those which form non-toxic base salts with the herein described acidic compounds of formula I. These non-toxic base salts include those derived from such pharmacologically acceptable cations as sodium, potassium, calcium and magnesium, etc. These salts can easily be prepared by treating the corresponding acidic compounds with an aqueous solution containing the desired pharmacologically acceptable cations, and then evaporating the resulting solution to dryness, preferably under reduced pressure. Alternatively, they may also be prepared by mixing lower alkanolic solutions of the acidic compounds and the desired alkali metal alkoxide together, and then evaporating the resulting solution to dryness in the same manner as before. In either case, stoichiometric quantities of reagents are preferably employed in order to ensure completeness of reaction and maximum product yields.

Compounds of the formula I and their pharmaceutically acceptable salts (hereinafter also referred to, collectively, as "the active compounds") are potent antagonists of the CCR1 receptors. The active compounds are useful in the treatment or prevention of autoimmune diseases (such as rheumatoid arthritis, type I diabetes (recent onset), inflammatory bowel disease, optic neuritis, psoriasis, multiple sclerosis, polymyalgia rheumatica, uveitis, and vasculitis), acute and chronic inflammatory conditions (such as osteoarthritis, adult respiratory distress syndrome, Respiratory Distress Syndrome of infancy, ischemia reperfusion injury, and glomerulonephritis), allergic conditions (such as asthma and atopic dermatitis), infection associated with inflammation (such as viral inflammation (including influenza and hepatitis) and Guillian-Barre), chronic bronchitis, xeno-transplantation, transplantation tissue rejection,



atherosclerosis, restenosis, HIV infectivity (co-receptor usage), and granulomatous diseases (including sarcoidosis, leprosy and tuberculosis).

The activity of the compounds of the invention can be assessed according to procedures known to those of ordinary skill in the art. Examples of recognized methods for determining CCR1 induced migration can be found in Coligan, J. E., Kruisbeek, A.M., Margulies, D.H., Shevach, E.M., Strober, W. editors: Current Protocols In Immunology, 6.12.1- 6.12.3. (John Wiley and Sons, NY, 1991). One specific example of how to determine the activity of a compound for inhibiting migration is described in detail below.

#### Chemotaxis Assay:

10       The ability of compounds to inhibit the chemotaxis to various chemokines can be evaluated using standard 48 or 96 well Boyden Chambers with a 5 micron polycarbonate filter. All reagents and cells can be prepared in standard RPMI (BioWhittaker Inc.) tissue culture medium supplemented with 1 mg/ml of bovine serum albumin. Briefly, MIP-1 $\alpha$  (Peprotech, Inc., P.O. Box 275, Rocky Hill NJ) or other test agonists, were placed into the  
15       lower chambers of the Boyden chamber. A polycarbonate filter was then applied and the upper chamber fastened. The amount of agonist chosen is that determined to give the maximal amount of chemotaxis in this system (e.g., 1 nM for MIP-1 $\alpha$  should be adequate).

THP-1 cells (ATCC TIB-202), primary human monocytes, or primary lymphocytes, isolated by standard techniques can then be added to the upper chambers in triplicate  
20       together with various concentrations of the test compound. Compound dilutions can be prepared using standard serological techniques and are mixed with cells prior to adding to the chamber.

After a suitable incubation period at 37 degrees centigrade (e.g. 3.5 hours for THP-1 cells, 90 minutes for primary monocytes), the chamber is removed, the cells in the upper  
25       chamber aspirated, the upper part of the filter wiped and the number of cells migrating can be determined according to the following method.

For THP-1 cells, the chamber (a 96 well variety manufactured by Neuroprobe) can be centrifuged to push cells off the lower chamber and the number of cells can be quantitated against a standard curve by a color change of the dye fluorescein diacetate.

30       For primary human monocytes, or lymphocytes, the filter can be stained with Dif Quik® dye (American Scientific Products) and the number of cells migrating can be determined microscopically.

The number of cells migrating in the presence of the compound are divided by the number of cells migrating in control wells (without the compound). The quotient is the %  
35       inhibition for the compound which can then be plotted using standard graphics techniques against the concentration of compound used. The 50% inhibition point is then determined

using a line fit analysis for all concentrations tested. The line fit for all data points must have an coefficient of correlation (R squared) of > 90% to be considered a valid assay.

All of the compounds of the invention that were tested had IC<sub>50</sub> of less than 25µM, in the Chemotaxis assay.

5           The compositions of the present invention may be formulated in a conventional manner using one or more pharmaceutically acceptable carriers. Thus, the active compounds of the invention may be formulated for oral, buccal, intranasal, parenteral (e.g., intravenous, intramuscular or subcutaneous) or rectal administration or in a form suitable for administration by inhalation or insufflation. The active compounds of the invention may also  
10 be formulated for sustained delivery.

For oral administration, the pharmaceutical compositions may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g., pregelatinized maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g., lactose, microcrystalline  
15 cellulose or calcium phosphate); lubricants (e.g., magnesium stearate, talc or silica); disintegrants (e.g., potato starch or sodium starch glycolate); or wetting agents (e.g., sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other  
20 suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g., sorbitol syrup, methyl cellulose or hydrogenated edible fats); emulsifying agents (e.g., lecithin or acacia); non-aqueous vehicles (e.g., almond oil, oily esters or ethyl alcohol); and preservatives (e.g., methyl or propyl p-hydroxybenzoates or sorbic acid).

25           For buccal administration, the composition may take the form of tablets or lozenges formulated in conventional manner.

The active compounds of the invention may be formulated for parenteral administration by injection, including using conventional catheterization techniques or infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampules or  
30 in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulating agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form for reconstitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

35           The active compounds of the invention may also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

For intranasal administration or administration by inhalation, the active compounds of the invention are conveniently delivered in the form of a solution or suspension from a pump spray container that is squeezed or pumped by the patient or as an aerosol spray presentation from a pressurized container or a nebulizer, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. The pressurized container or nebulizer may contain a solution or suspension of the active compound. Capsules and cartridges (made, for example, from gelatin) for use in an inhaler or insufflator may be formulated containing a powder mix of a compound of the invention and a suitable powder base such as lactose or starch.

A proposed dose of the active compounds of the invention for oral, parenteral or buccal administration to the average adult human for the treatment of the conditions referred to above (e.g., rheumatoid arthritis) is 0.1 to 1000 mg of the active ingredient per unit dose which could be administered, for example, 1 to 4 times per day.

Aerosol formulations for treatment of the conditions referred to above (e.g., rheumatoid arthritis) in the average adult human are preferably arranged so that each metered dose or "puff" of aerosol contains 20  $\mu$ g to 1000  $\mu$ g of the compound of the invention. The overall daily dose with an aerosol will be within the range 0.1 mg to 1000 mg. Administration may be several times daily, for example 2, 3, 4 or 8 times, giving for example, 1, 2 or 3 doses each time.


The active agents can be formulated for sustained delivery according to methods well known to those of ordinary skill in the art. Examples of such formulations can be found in United States Patents 3,538,214, 4,060,598, 4,173,626, 3,119,742, and 3,492,397.

The compounds of the invention can also be utilized in combination therapy with, but not limited to, other therapeutic agents such as with T-cell immunosuppressant agents such as rapamycin cyclosporin A and FK-506, with steroid sparing agents such as Cellcept®, or with classical anti-inflammatory agents (e.g. cyclooxygenase/lipoxygenase inhibitors) such as tenidap, aspirin, acetaminophen, naproxen and piroxicam.

The following Examples illustrate the preparation of the compounds of the present invention. NMR data are reported in parts per million ( $\delta$ ) and are referenced to the deuterium lock signal from the sample solvent (deuteriochloroform unless otherwise specified). Commercial reagents were utilized without further purification. THF refers to tetrahydrofuran. DMF refers to N,N-dimethylformamide. Chromatography refers to column chromatography performed using 32-63 mm silica gel and executed under nitrogen pressure (flash chromatography) conditions. Low Resolution Mass Spectra (LRMS) were recorded on either a Hewlett Packard 5989®, utilizing chemical ionization (ammonium), or a Fisons (or Micro

Mass) Atmospheric Pressure Chemical Ionization (APCI) platform which uses a 50/50 mixture of acetonitrile/water with 0.1% formic acid as the ionizing agent. Room or ambient temperature refers to 20-25°C. All non-aqueous reactions were run under a nitrogen atmosphere for convenience and to maximize yields. The names for the compounds of the invention were created by the Autonom 2.0 PC-batch version from Bellstein Informationssysteme GmbH (ISBN 3-89536-976-4).

### Example 1



### (2R,5S)-2-[4-Chloro-2-(piperazine-1-carbonyl)-phenoxy]-1-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-ethanone

#### (R)-2-(4-Fluoro-benzylamino)-propionic acid methyl ester

To a solution of (R)-2-amino-propionic acid methyl ester hydrochloride (25 g, 179 mmol) and 4-fluorobenzaldehyde (23 ml, 215 mmol) in 1,2-dichloroethane (200 ml) was added triethylamine (25 ml, 179 mmol). The resulting mixture was stirred for two hours at ambient temperature followed by addition of sodium acetoxyborohydride (57 g, 268 mmol) in four portions. The resulting mixture was stirred overnight at ambient temperature. The reaction was neutralized with dilute aqueous sodium hydroxide solution and extracted with dichloromethane (2X). The organic layers were combined, dried over magnesium sulfate, filtered and concentrated in vacuo. Chromatography on silica gel gave the title compound (34.4 g, 91% yield).

#### (2R,5S)-2-[(2-tert-Butoxycarbonylamino-propionyl)-(4-fluoro-benzyl)-amino]-propionic acid methyl ester

To a solution of (R)-2-tert-butoxycarbonylamino-propionic acid (37 g, 195 mmol) in dry tetrahydrofuran (250 ml) at 0 °C was added 4-methyl morpholine (21.5 ml, 195 mmol) followed by isobutylchloroformate (25.3 ml, 195 mmol). The reaction was allowed to warm to ambient temperature and stirred for two hours. This was followed by the addition of (S)-2-(4-fluoro-benzylamino)-propionic acid methyl ester (34.4 g, 162 mmol). The resulting mixture was stirred overnight at ambient temperature. The reaction mixture was filtered through a pad of celite and the filter cake was washed with ethyl acetate. The filtrate was concentrated in vacuo, diluted with ethyl acetate and washed with water and brine. The organics were

dried over magnesium sulfate, filtered and concentrated in vacuo. Chromatography on silica gel gave the title compound (43.2 g, 70% yield).

(2R,5S)-1-(4-Fluoro-benzyl)-3,6-dimethyl-piperazine-2,5-dione

To a solution of (2R,5S)-2-[(2-tert-butoxycarbonylamino-propionyl)-(4-fluoro-benzyl)-amino]-propionic acid methyl ester (43 g, 382 mmol) in dichloromethane (120 ml) at 0 °C was added trifluoroacetic acid (60 ml). The reaction was allowed to warm to ambient temperature and stirred for 2 hours. The reaction was cooled to 0 °C and slowly quenched by addition of 3N aqueous sodium hydroxide until basic. The resulting mixture was extracted with dichloromethane (2X). The combined organics were dried over magnesium sulfate, filtered and concentrated in vacuo to give the title compound (22 g, 78% yield).

(2R, 5S)-1-(4-Fluoro-benzyl)-2,5-dimethyl-piperazine

To a solution of (2R, 5S)-1-(4-fluoro-benzyl)-3,6-dimethyl-piperazine-2,5-dione (22 g, 87.9 mmol) in dry tetrahydrofuran (160 ml) at 0 °C was added a solution of lithium aluminum hydride (1M in tetrahydrofuran, 373 ml, 373 mmol) dropwise over 40 minutes. The reaction mixture was then refluxed for 4 hours, cooled to ambient temperature and slowly quenched with water. The resulting mixture was filtered through a pad of celite and the filter cake was washed with ethyl acetate. The filtrate was then concentrated, diluted with ethyl acetate and washed with saturated aqueous sodium hydrogen carbonate. The organic layer was separated, dried over magnesium sulfate, filtered and concentrated in vacuo to give the title compound (17.7 g, 91% yield).

(2R, 5S)-2-Chloro-1-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-ethanone

To a solution of (2R, 5S)-1-(4-fluoro-benzyl)-2,5-dimethyl-piperazine (2.5 g, 11.25mmol) in dry dichloromethane (11 ml) at 0 °C was added triethylamine (1.57 ml, 11.2 mmol) followed by chloroacetyl chloride (0.858 ml, 11.2 mmol). The resulting reaction mixture was stirred for 30 minutes. The reaction was then filtered through a pad of celite, washed with dichloromethane and the resulting filtrate was concentrated to give a yellow oil. Chromatography on silica gel gave the title compound (2.84 g, 86% yield).

(2R, 5S)-5-Chloro-2-[2-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-2-oxo-ethoxy]-benzoic acid methyl ester

To a solution of (2R, 5S)-2-chloro-1-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-ethanone (2.75 g, 9.2 mmol) in butanone (50 ml) was added methyl 5-chloro-2-hydroxybenzoate (1.55 g, 9.2 mmol), potassium carbonate (2.54 g, 18.4 mmol) and potassium iodide (1.52 g, 9.2 mmol). The resulting mixture was stirred at reflux for 6 hours. The reaction was then cooled, diluted with ethyl acetate, and washed with brine. The organics were dried over magnesium sulfate, filtered and concentrated in vacuo to give an orange oil. Chromatography on silica gel gave the title compound (4.1 g, 100% yield).

(2R, 5S)-5-Chloro-2-[2-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-2-oxo-ethoxy]-benzoic acid

To a solution of (2R, 5S)-5-chloro-2-[2-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-2-oxo-ethoxy]-benzoic acid methyl ester (4.12 g, 9.18 mmol) in tetrahydrofuran (10 ml), methanol (10 ml) and water (4 ml) was added lithium hydroxide monohydrate (1.93 g, 45.9 mmol). The resulting mixture was stirred overnight at ambient temperature. The reaction was then concentrated, diluted with 1N hydrochloric acid and extracted with dichloromethane (2X). The organic layers were combined, dried over magnesium sulfate, filtered and concentrated to give a white foam. Triteration in dichloromethane and diethyl ether gave the title compound (1.38 g, 35% yield).


(2R, 5S)-4-(5-Chloro-2-[2-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-2-oxo-ethoxy]-benzoyl)-piperazine-1-carboxylic acid tert-butyl ester

To a solution of (2R, 5S)-5-chloro-2-[2-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-2-oxo-ethoxy]-benzoic acid (0.10 g, 0.212 mmol) in dichloromethane (4 ml) was added 4-dimethylaminopyridine (0.039 g, 0.318 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.061 g, 0.318 mmol) and *tert*-butyl 1-piperazinecarboxylate (0.041 g, 0.222 mmol). The resulting mixture was stirred overnight at ambient temperature. It was then diluted with dichloromethane and washed with brine. The organics were dried over magnesium sulfate, filtered and concentrated to give a clear oil. Chromatography on silica gel gave the title compound (0.110 g, 85% yield).


(2R, 5S)-2-[4-Chloro-2-(piperazine-1-carbonyl)-phenoxy]-1-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-ethanone

To a solution of (2R, 5S)-4-(5-chloro-2-[2-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-2-oxo-ethoxy]-benzoyl)-piperazine-1-carboxylic acid *tert*-butyl ester (0.110 g, 0.182 mmol) in dichloroethane (10 ml) was added trifluoroacetic acid (5 ml). The reaction was stirred for two hours at ambient temperature. The reaction was then diluted with dichloromethane and washed with 1N aqueous sodium hydroxide. The organics were dried over magnesium sulfate, filtered and concentrated to give the title compound (0.080 g, 87% yield).

The title compounds for Examples 2 – 12 were prepared by a method analogous to that described in Example 1.



Example	R <sup>20</sup>	R <sup>2</sup>
2		Me
3		Me
4		H
5		H
6		H
7		H
8		H
9		H
10	-NHSO <sub>2</sub> Me	Me
11	-NH-(CH <sub>2</sub> ) <sub>2</sub> -NHSO <sub>2</sub> CH <sub>3</sub>	CH <sub>3</sub>
12	-NH-(CH <sub>2</sub> ) <sub>2</sub> -NHC(O)NH <sub>2</sub>	CH <sub>3</sub>

**Example 13**

**(2R)-3-Amino-N-(5-chloro-2-{2-[4-(4-fluoro-benzyl)-2-methyl-piperazin-1-yl]-2-oxo-ethoxy}-phenyl)-propionamide**

5        **[2-(5-Chloro-2-{2-[4-(4-fluoro-benzyl)-2-methyl-piperazin-1-yl]-2-oxo-ethoxy}-phenylcarbamoyl)-ethyl]-carbamic acid tert-butyl ester**


To a solution of 2-(2-amino-4-chloro-phenoxy)-1-[4-(4-fluoro-benzyl)-2-methyl-piperazin-1-yl]-ethanone (0.066 g, 0.17 mmol) in methylene chloride (2 mL) at ambient temperature was added N-methylmorpholine (0.025 mL, 0.23 mmol), 3-tert-butoxycarbonylamino-propionic acid (0.044 g, 0.23 mmol) and O-benzotriazole-1-yl-N,N,N',N'-tetramethyluronium hexafluorophosphate (0.076 g, 0.20 mmol). The resulting solution was stirred at ambient temperature for 60 hours, then concentrated. Radial chromatography (2 mm plate) gave the title compound (0.114 g)

15        **(2R)-3-Amino-N-(5-chloro-2-{2-[4-(4-fluoro-benzyl)-2-methyl-piperazin-1-yl]-2-oxo-ethoxy}-phenyl)-propionamide**


To a solution of [2-(5-chloro-2-{2-[4-(4-fluoro-benzyl)-2-methyl-piperazin-1-yl]-2-oxo-ethoxy}-phenylcarbamoyl)-ethyl]-carbamic acid tert-butyl ester (0.110 g, 0.2 mmol) in methylene chloride (3 mL) was added trifluoroacetic acid (0.50 mL). The reaction was stirred for 2 hours at ambient temperature then diluted with saturated aqueous sodium hydrogen carbonate. The mixture was extracted with methylene chloride and the combined organics were dried over magnesium sulfate, filtered and concentrated in vacuo to give the title compound (0.069 g)

20        The title compounds for Examples 14 – 19 were prepared by a method analogous to that described in Example 13.





Example	R <sup>20</sup>	R <sup>2</sup>
14		H
15		H
16		H
17		H
18		H
19		H

**Example 20**

- 5 (2R,5S)-1-(2-Amino-ethyl)-3-(5-chloro-2-{2-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-2-oxo-ethoxy}-phenyl)-urea

(2R, 5S)-2-(4-Chloro-2-nitro-phenoxy)-1-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-ethanone

To a solution of (2R, 5S)-5-chloro-2-{2-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-2-oxo-ethoxy}-benzoic acid methyl ester (1.0 g, 3.35 mmol) in butanone (35 ml) was added  
5 2-nitro-4-chlorophenol (0.639 g, 3.69 mmol), potassium carbonate (0.925 g, 6.7mmol) and potassium iodide (0.556 g, 3.35 mmol). The reaction mixture was heated at reflux overnight. The reaction mixture was then cooled, diluted with water and extracted with ethyl acetate. The combined organics were dried over magnesium sulfate, filtered and concentrated to give an orange oil. Chromatography on silica gel gave the title compound (1.35 g, 93% yield).

10 (2R, 5S)-2-(2-Amino-4-chloro-phenoxy)-1-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-ethanone

To a solution of (2R, 5S)-2-(4-chloro-2-nitro-phenoxy)-1-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-ethanone (2.2 g, 5.05 mmol) in ethanol (50 ml) in a par bottle was added 5% platinum on carbon (2.2 g). The reaction mixture was subjected to hydrogen gas  
15 (35 psi) for thirty minutes. The reaction mixture was filtered through celite and washed with ethanol. The filtrate was concentrated to give a tan foam. Chromatography on silica gel gave the title compound (1.42 g, 70% yield).


(2R, 5S)-(5-Chloro-2-{2-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-2-oxo-ethoxy}-phenyl)-carbamic acid 4-nitro-phenyl ester

20 To a solution of (2R, 5S)-2-(2-amino-4-chloro-phenoxy)-1-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-ethanone (0.150 g, 0.37 mmol) in dichloromethane (7 ml) was added pyridine (0.066 ml, 0.82 mmol) followed by 4-nitrophenyl chloroformate (0.075 g, 0.41 mmol). The reaction was stirred at ambient temperature for 3 1/2 hours. The reaction mixture was concentrated followed by chromatography on silica gel to give the title compound (0.153 g,  
25 74%yield).


(2R,5S)-1-(2-Amino-ethyl)-3-(5-chloro-2-{2-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-2-oxo-ethoxy}-phenyl)-urea

To a solution of (2R, 5S)-(5-chloro-2-{2-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-2-oxo-ethoxy}-phenyl)-carbamic acid 4-nitro-phenyl ester (0.206 g, 0.37 mmol) in dry  
30 methanol (6 ml) was added ethyldiamine (0.05 ml, 0.814 mmol). The reaction was stirred at ambient temperature overnight. The reaction was concentrated and chromatographed on silica gel to give the title compound (0.115 g, 63% yield).

The title compounds for Examples 21 – 27 were prepared by a method analogous to that described in Example 20.



Example	R <sup>20</sup>	R <sup>2</sup>
21		Me
22		H
23		Me
24		H
25		H
26		H
27		Me

**Example 28**

**(2R)-2-Amino-ethanesulfonic acid (5-chloro-2-{2-[4-(4-fluoro-benzyl)-2-methyl-piperazin-1-yl]-2-oxo-ethoxy}-phenyl)-amide**


**2-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-ethanesulfonic acid (5-chloro-2-{2-[4-(4-fluoro-benzyl)-2-methyl-piperazin-1-yl]-2-oxo-ethoxy}-phenyl)-amide**





5 To a solution of of 2-(2-amino-4-chloro-phenoxy)-1-[4-(4-fluoro-benzyl)-2-methyl-piperazin-1-yl]-ethanone (0.050 g, 0.13 mmol) in methylene chloride (1 mL) at ambient temperature was added triethylamine (0.027 mL, 0.19 mmol) and 2-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-ethanesulfonyl chloride (0.045 g, 0.17 mmol). The reaction was stirred overnight at ambient temperature. Additional triethylamine ((0.027 mL, 0.19 mmol) and 2-  
10 (1,3-dioxo-1,3-dihydro-isoindol-2-yl)-ethanesulfonyl chloride (0.045 g, 0.17 mmol) was added. The reaction was stirred one hour, then additional triethylamine (0.055 mL, 0.34 mmol) was added. The reaction was stirred and hour, then additional 2-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-ethanesulfonyl chloride (0.090 g, 0.34 mmol) was added. After stirring an additional 1  
15 hour, the reaction was treated with saturated aqueous sodium hydrogen carbonate and extracted with methylene chloride (3X). The combined organics were dried over magnesium sulfate, filtered and concentrated in vacuo. Purification via radial chromatography (2 mm plate) gave the title compound (0.030 g).


**(2R)-2-Amino-ethanesulfonic acid (5-chloro-2-{2-[4-(4-fluoro-benzyl)-2-methyl-piperazin-1-yl]-2-oxo-ethoxy}-phenyl)-amide**

20 To a solution of 2-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-ethanesulfonic acid (5-chloro-2-{2-[4-(4-fluoro-benzyl)-2-methyl-piperazin-1-yl]-2-oxo-ethoxy}-phenyl)-amide (0.030 g, 0.048 mmol) in EtOH (1 mL) at ambient temperature was added hydrazine hydrate (0.025 mL). The reaction was stirred overnight at ambient temperature then diluted with water and extracted with methylene chloride (2X). The combined organics were washed with saturated aqueous  
25 brine and dried over sodium sulfate, filtered and concentrated in vacuo. Purification via radial chromatography (2 mm plate) gave the title compound (0.014 g).

The title compounds for Examples 29 – 34 were prepared by a method analogous to that described in Example 28.



Example	R <sup>20</sup>	R <sup>2</sup>
29	-SO <sub>2</sub> CF <sub>3</sub>	Me
30		H
31		H
32		H
33		H
34	-SO <sub>2</sub> N(Me) <sub>2</sub>	Me

**Example 35**

**(2R)-N-(5-Chloro-2-{2-[4-(4-fluorobenzyl)-2-methylpiperazin-1-yl]-2-oxo-ethoxy}-phenyl)-cyanoguanidine**



**1-(5-Chloro-2-{2-[4-(4-fluorobenzyl)-2-methylpiperazin-1-yl]-2-oxo-ethoxy}-phenyl)-N-cyano-S-methyl-isothiurea**

To a solution of 2-(2-amino-4-chlorophenoxy)-1-[4-(4-fluorobenzyl)-2-methylpiperazin-1-yl]ethanone (0.30 g, 0.77 mmol) in tetrahydrofuran (5 mL) at ambient temperature was added sodium hydride (0.029 g, 1.22 mmol) and the reaction was stirred for 30 minutes. To this was added S,S1-dimethyl N-cyanodithio iminocarbonate (0.168, 1.15 mmol) and the mixture was heated at reflux overnight. The reaction was cooled and quenched with saturated aqueous ammonium chloride. The mixture was extracted with EtOAc and the combined organics were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated in vacuo. Chromatography on silica gel gave the title compound (0.350 g)

(2R)-N-(5-Chloro-2-{2-[4-(4-fluoro-benzyl)-2-methyl-piperazin-1-yl]-2-oxo-ethoxy}-phenyl)-N-cyano-S-methyl-isothiurea


- 5 To a solution of 1-(5-chloro-2-{2-[4-(4-fluoro-benzyl)-2-methyl-piperazin-1-yl]-2-oxo-ethoxy}-phenyl)-N-cyano-S-methyl-isothiurea (0.045 g, 0.092 mmol) in EtOH (1 mL) was added ammonium hydroxide (0.100 mL) and the resulting solution was shaken at 60 °C overnight. The crude reaction mixture was purified directly via radial chromatography (2 mm plate) to give the title compound (0.027 g).

The title compounds for Examples 36 -- 38 were prepared by a method analogous to that described in Example 35.



10

Example	R <sup>20</sup>	R <sup>2</sup>
36		H
37		H
38		H

**Example 39**

**(2R,5S)-2-{4-Chloro-2-[(2-diethylamino-ethylamino)-methyl]-phenoxy}-1-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-ethanone**

5 **(2R,5S)-5-Chloro-2-[2-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-2-oxo-ethoxy]-benzaldehyde**


To a solution of 2-chloro-1-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-ethanone (2.87 g, 9.6 mmol) in DMF (20 mL) was added 5-chlorosalicylaldehyde (1.65 g, 10.5 mmol), potassium carbonate (2.64 g, 19.2 mmol) and potassium iodide (1.59 g, 9.6 mmol). The resulting mixture was heated to 100 °C for 12 hours. The reaction was cooled, diluted with saturated aqueous brine and extracted with ethyl acetate (3X). The combined organics were dried over magnesium sulfate and filtered. The filtrate was concentrated in vacuo to give crude product. Purification via chromatography on silica gel (15% EtOAc/Hexanes) gave the title compound 3.40 g, 85% yield.)

15 **(2R,5S)-2-{4-Chloro-2-[(2-diethylamino-ethylamino)-methyl]-phenoxy}-1-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-ethanone**

To a solution of (2R,5S)-5-chloro-2-[2-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-2-oxo-ethoxy]-benzaldehyde (0.100 g, 0.25 mmol) in 10:1 dichloroethane acetic acid (2.2 mL) was added (diethylamino)ethylamine (0.088 mL, 0.625 mmol) and the resulting solution was stirred for 1 hour at ambient temperature. To this was added sodium cyanoborohydride (0.0094 g, 0.15 mmol) and the reaction was stirred overnight at ambient temperature. Upon completion water was added and the mixture was basified with solid sodium bicarbonate (pH > 10). The product was extracted with dichloromethane (2X) and diethyl ether (2X). The combined organics were dried over magnesium sulfate, filtered and concentrated in vacuo. Chromatography on silica gel gave the title compound (0.039 g, x 30% yield)

The title compounds for Examples 40 – 62 were prepared by a method analogous to that described in Example 39.


-55-



Example	R <sup>20</sup>	R <sup>2</sup>
40		Me
41		Me
42		Me
43		Me
44		Me
45		Me
46		Me
47		Me
48		Me
49		Me
50		Me
51		Me



Example	R <sup>20</sup>	R <sup>2</sup>
52		Me
53		Me
54		Me
55		Me
56		Me
57		Me
58		Me
59		Me
60		Me
61		Me
62		Me

**Example 63**

**N-(5-Chloro-2-{2-[4-(4-fluorobenzyl)-2,5-dimethylpiperazin-1-yl]-2-oxoethoxy}-benzyl)-2-diethylaminoacetamide**

**(2R,5S)-2-(2-Aminomethyl-4-chlorophenoxy)-1-[4-(4-fluorobenzyl)-2,5-dimethylpiperazin-1-yl]ethanone**


To a solution of (2R,5S)-5-chloro-2-[2-[4-(4-fluorobenzyl)-2,5-dimethylpiperazin-1-yl]-2-oxoethoxy]benzaldehyde (1.86 g, 4.44 mmol) in MeOH (20 mL) was added ammonium

acetate (3.42 g, 44 mmol) and sodium cyanoborohydride (0.195 g, 3.1 mmol). The resulting mixture was stirred overnight at ambient temperature. The reaction was quenched with concentrated hydrochloric acid and concentrated in vacuo. The residue was dissolved in water and basified with aqueous 3N NaOH (pH > 10). The product was extracted with dichloromethane (2X) and ethyl acetate (2X). The combined organics were dried over magnesium sulfate, filtered and concentrated in vacuo. Chromatography on silica gel gave the title compound (1.29 g, 69 % yield)

N-(5-Chloro-2-[2-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-2-oxo-ethoxy]-benzyl)-2-diethylamino-acetamide















To a solution of N,N-dimethylglycine (0.014 g, 0.13 mmol) and (2R,5S)-2-(2-aminomethyl-4-chloro-phenoxy)-1-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-ethanone (0.063 g, 0.15 mmol) in dichloromethane (2 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.034 g, 0.18 mmol), 1-hydroxybenzotriazole (0.021g, 0.15 mmol), and triethylamine (0.036 mL, 0.36 mmol). After the reaction was stirred for 48 hours, the solution was diluted with saturated aqueous sodium bicarbonate and extracted with dichloromethane (3X). The combined organics were dried over magnesium sulfate, filtered and concentrated in vacuo. Chromatography on silica gel gave the title compound (0.063 g, 80 %).






The title compounds for Examples 64 – 85 were prepared by a method analogous to that described in Example 63.




20

Example	R <sup>20</sup>	R <sup>2</sup>
64		Me
65		Me
66		Me

Example	R <sup>20</sup>	R <sup>2</sup>
67		Me
68		Me
69		Me
70		Me
71		Me
72		Me
73		Me
74		Me
75		Me
76		Me
77		Me
78		H
79		H
80		Me

Example	R <sup>20</sup>	R <sup>2</sup>
81		Me
82		Me
83		Me
84		H
85		H

**Example 86**

**(2R,5S)-(N-{2-[3-(5-Chloro-2-{2-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-2-oxo-ethoxy}-benzyl)-ureido]-ethyl}-acetamide**

5


To a solution of (2R,5S)-5-chloro-2-{2-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-2-oxo-ethoxy}-benzylamine (0.200 g, 0.477 mmol) in methylene chloride (10 mL) was added pyridine (0.077 mL, 0.954 mmol) and 4-nitro phenyl chloroformate (0.097 g, 0.525 mmol). The resulting mixture was stirred for one hour at ambient temperature and then concentrated in vacuo. The residue (0.055g, 0.094 mmol) was dissolved in methanol (1 mL). N-Acetyleneethylenediamine (0.019 mL, 0.188 mmol) was added and the reaction was stirred at room temperature over night. The reaction was concentrated in vacuo and chromatography on silica gel gave the title compound (0.019 g, 27 %).

10


The title compounds for Examples 87 – 90 were prepared by a method analogous to that described in Example 86.

15

-60-



Example	R <sup>20</sup>	R <sup>2</sup>
87		Me
88		Me
89		Me
90		Me

**Example 91**

5

**(2R,5S)- 2-Amino-ethanesulfonic acid 5-chloro-2-(2-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-2-oxo-ethoxy)-benzylamide**

**(2R,5S)-2-(1,3-Dioxo-1,3-dihydro-isolindol-2-yl)-ethanesulfonic acid 5-chloro-2-(2-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-2-oxo-ethoxy)-benzylamide**


10 To a solution of (2R,5S)-5-chloro-2-(2-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-2-oxo-ethoxy)-benzyl amine (0.060 g, 0.143 mmol) in methylene chloride (3 mL) was added 2-(1,3-dioxo-1,3-dihydro-isolindol-2-yl)-ethanesulfonyl chloride (0.043 g, 0.150 mmol) and triethylamine (0.060 mL, 0.43 mmol) and the solution was stirred 1 hr at ambient

temperature. The reaction was diluted with saturated aqueous sodium hydrogen carbonate and extracted with methylene chloride. The combined organics were dried over magnesium sulfate, filtered and concentrated in vacuo. Chromatography on silica gel gave the title compound (0.073 g, 77%).

5      (2R,5S)- 2-Amino-ethanesulfonic acid 5-chloro-2-{2-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-2-oxo-ethoxy}-benzylamide


To a solution of (2R,5S)-2-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-ethanesulfonic acid 5-chloro-2-{2-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-2-oxo-ethoxy}-benzylamide (0.073 g, 0.111 mmol) in EtOH (1 mL) was added hydrazine (35% aq. 0.25 mL, 2.73 mmol) and the solution was stirred overnight at ambient temperature. The reaction was filtered through a glass frit and washed with EtOH. The filtrate was concentrated in vacuo to give the title compound (0.056 g, 96%)

The title compounds for Examples 92 – 93 were prepared by a method analogous to that described in Example 91.




15

Example	R <sup>20</sup>	R <sup>2</sup>
92	$\text{-SO}_2\text{-CH}_2\text{-NH}_2$	H
93	$\text{-SO}_2\text{NMe}_2$	Me


**Example 94**

**(2R)-N-(2-Amino-ethyl)-N'-(5-chloro-2-{2-[4-(4-fluoro-benzyl)-2-methyl-piperazin-1-yl]-2-oxo-ethoxy}-benzyl)-cyanoguanidine**

- 5 A solution of 2-(2-aminomethyl-4-chloro-phenoxy)-1-[4-(4-fluoro-benzyl)-2-methyl-piperazin-1-yl]-ethanone (0.025 g, 0.062 mmol) and diphenyl cyanocarbonimidate (0.016 g, 0.068 mmol) in ethanol (1 mL) was heated on a shaker plate at 60°C. After 22 h, ethylenediamine (0.008 mL, 0.123 mmol) was added and the resulting solution was heated on a shaker plate at 60°C for an additional 21 h. The solution was cooled to ambient
- 10 temperature, concentrated and purified using radial chromatography to yield the title compound (0.021 g, 67%). The title compounds for Examples 95 – 96 were prepared by a method analogous to that described in Example 94.



Example	R <sup>20</sup>	R <sup>2</sup>
95		H
96		H

**Example 97**

**(2R, 5S)-5-Chloro-2-{2-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-2-oxo-ethoxy}-N-methyl-benzenesulfonamide**

**5      5-Chloro-2-methoxy-N-methyl-benzenesulfonamide**

To a solution of 5-Chloro-2-methoxy-benzenesulfonyl chloride (1.0g, 4.15 mmol) in tetrahydrofuran(8ml) was bubbled methylamine gas until saturated. The reaction mixture was sealed with a septa and stirred overnight at ambient temperature. The reaction mixture was then concentrated, triterated in dichloromethane and ether, filtered and dried yielding the above titled compound 1.05 g (>100%) white solid.

**5-Chloro-2-hydroxy-N-methyl-benzenesulfonamide**


To a solution of sodium hydride (60% in mineral oil, 90.24 mg, 2.25mmol) in dry dimethyl formamide was added thiophenol (0.225 ml, 2.25mmol) dropwise. To this was then added 5-Chloro-2-methoxy-N-methyl-benzenesulfonamide (531mg, 2.25mmol) followed by refluxing for 4 hours. The reaction mixture was cooled, diluted with ethyl acetate and washed with a 1N sulfuric acid solution. The organic layer was separated, dried over magnesium sulfate, filtered and concentrated in vacuo. Chromatography on silica gel yielded the above titled compound (320mg, 53% yield).

**20      (2R, 5S)-5-Chloro-2-{2-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-2-oxo-ethoxy}-N-methyl-benzenesulfonamide**

To a solution of (2R, 5S)-2-Chloro-1-[4-(4-fluoro-benzyl)-2,5-dimethyl-piperazin-1-yl]-ethanone (375mg, 1.26 mmol) in butanone (12ml) was added 5-Chloro-2-hydroxy-N-methyl-benzenesulfonamide (280mg, 1.26mmol), potassium carbonate (348mg, 2.52mmol) and potassium iodide (209mg, 1.26mmol). The resulting reaction mixture was refluxed for 4 hours. It was allowed to cool, diluted with ethyl acetate and washed with brine. The organic layer was separated, dried over magnesium sulfate, filtered and concentrated in vacuo. Chromatography on silica gel gave the above titled compound (320mg, 53% yield).




The title compounds for Examples 98 – 100 were prepared by a method analogous to that described in Example 97.



Example	R <sup>20</sup>	R <sup>2</sup>
98	-NH <sub>2</sub>	Me
99		Me
100		Me

## CLAIMS FOR FOREIGN FILING

1. A method of preparing a compound of the formula



or the pharmaceutically acceptable salt thereof; wherein

- 5           a is 1, 2, 3, 4 or 5;  
             b is 0, 1, 2, 3 or 4;  
             c is 0 or 1;  
             d is 1, 2, 3, 4 or 5;  
             e is 0 or 1;
- 10           j is 1, 2, 3, or 4;  
             X is C(O), C(S) or CH<sub>2</sub>;  
             Y is CH<sub>2</sub>, or if e is 0, Y is CHR<sup>8</sup> wherein R<sup>8</sup> is hydrogen, (C<sub>6</sub>-C<sub>10</sub>)aryl or NR<sup>9</sup>R<sup>10</sup>;  
             Z is oxygen, NR<sup>9</sup> or CR<sup>11</sup>R<sup>12</sup>;
- each R<sup>1</sup> is independently selected from hydrogen, hydroxy, hydroxysulfonyl, halo,  
 15 (C<sub>1</sub>-C<sub>6</sub>)alkyl, mercapto, mercapto(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylthio, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylthio(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>6</sub>-C<sub>10</sub>)aryloxy, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, trifluoromethyl, formyl, formyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, nitro, nitroso, cyano, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, trifluoromethoxy, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkylamino, (C<sub>3</sub>-C<sub>7</sub>)cycloalkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>3</sub>-C<sub>7</sub>)cycloalkyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, ((C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyano(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>7</sub>)alkenyl, (C<sub>2</sub>-C<sub>7</sub>)alkynyl, (C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylthio(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino, (C<sub>6</sub>-C<sub>10</sub>)arylamino, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, carboxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-

$C_{10}$ arylcarbonyl,  $(C_8-C_{10})$ arylcarbonyl $(C_1-C_6)$ alkyl,  $(C_8-C_{10})$ aryl $(C_1-C_6)$ alkylcarbonyl,  $(C_8-C_{10})$ aryl $(C_1-C_6)$ alkylcarbonyl $(C_1-C_6)$ alkyl, carboxy $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxycarbonyl $(C_1-C_6)$ alkyl,  $(C_8-C_{10})$ aryl $(C_1-C_6)$ alkoxycarbonyl $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxy $(C_1-C_6)$ alkylcarbonyloxy $(C_1-C_6)$ alkyl, aminocarbonyl,  $(C_1-C_6)$ alkylaminocarbonyl,  $((C_1-C_6)$ alkyl) $_2$ aminocarbonyl,  $(C_8-C_{10})$ arylaminocarbonyl,  $(C_8-C_{10})$ aryl $(C_1-C_6)$ alkylaminocarbonyl, aminocarbonyl $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylaminocarbonyl $(C_1-C_6)$ alkyl,  $((C_1-C_6)$ alkyl) $_2$ aminocarbonyl $(C_1-C_6)$ alkyl,  $(C_8-C_{10})$ arylaminocarbonyl $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylaminocarbonyl $(C_1-C_6)$ alkyl, amidino, guanidino, ureido,  $(C_1-C_6)$ alkylureido,  $((C_1-C_6)$ alkyl) $_2$ ureido, ureido $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylureido $(C_1-C_6)$ alkyl,  $((C_1-C_6)$ alkyl) $_2$ ureido $(C_1-C_6)$ alkyl,  $(C_2-C_9)$ heterocycloalkyl,  $(C_2-C_9)$ heteroaryl,  $(C_2-C_9)$ heterocycloalkyl $(C_1-C_6)$ alkyl and  $(C_2-C_9)$ heteroaryl $(C_1-C_6)$ alkyl;

each  $R^2$  and  $R^3$  are independently selected from oxo, halo,  $(C_1-C_6)$ alkyl,  $(C_3-C_8)$ cycloalkyl,  $(C_3-C_8)$ cycloalkyl $(C_1-C_6)$ alkyl,  $(C_3-C_8)$ cycloalkylamino $(C_1-C_6)$ alkyl,  $(C_3-C_8)$ cycloalkyl $(C_1-C_6)$ alkylamino $(C_1-C_6)$ alkyl, halo $(C_1-C_6)$ alkyl,  $(C_2-C_6)$ alkenyl,  $(C_2-C_6)$ alkynyl,  $(C_8-C_{10})$ aryl,  $(C_8-C_{10})$ aryl $(C_1-C_6)$ alkyl,  $(C_8-C_{10})$ aryl $(C_2-C_6)$ alkenyl, H-C(O)-, H-C(O)-(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy $(C_1-C_6)$ alkyl, hydroxy $(C_2-C_6)$ alkenyl, hydroxy $(C_2-C_6)$ alkynyl, hydroxy $(C_8-C_{10})$ aryl $(C_1-C_6)$ alkyl, hydroxy $(C_3-C_8)$ cycloalkyl $(C_1-C_6)$ alkyl, thio $(C_1-C_6)$ alkyl, cyano $(C_1-C_6)$ alkyl, halo $(C_1-C_6)$ alkylcarbonylamino $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxy $(C_8-C_{10})$ aryl $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxy $(C_1-C_6)$ alkyl,  $(C_8-C_{10})$ aryloxy $(C_1-C_6)$ alkyl,  $(C_8-C_{10})$ aryl $(C_1-C_6)$ alkoxy $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylthio $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylsulfinyl $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylsulfonyl $(C_1-C_6)$ alkyl, hydroxy $(C_1-C_6)$ alkylthio $(C_1-C_6)$ alkyl, amino $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylamino $(C_1-C_6)$ alkyl,  $((C_1-C_6)$ alkyl) $_2$ amino $(C_1-C_6)$ alkyl,  $(C_8-C_{10})$ arylamino $(C_1-C_6)$ alkyl,  $(C_8-C_{10})$ aryl $(C_1-C_6)$ alkylamino $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylcarbonylamino $(C_1-C_6)$ alkyl, azido $(C_1-C_6)$ alkyl, aminocarbonylamino $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylaminocarbonylamino $(C_1-C_6)$ alkyl,  $((C_1-C_6)$ alkyl) $_2$ aminocarbonylamino $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxycarbonyl $(C_1-C_6)$ alkylaminocarbonylamino $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxycarbonylamino $(C_1-C_6)$ alkyl, hydroxy $(C_1-C_6)$ alkylamino $(C_1-C_6)$ alkyl,  $(C_8-C_{10})$ aryloxy $(C_1-C_6)$ alkylcarbonyloxy $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxy $(C_1-C_6)$ alkylcarbonyloxy $(C_1-C_6)$ alkyl,  $(C_8-C_{10})$ aryl $(C_1-C_6)$ alkoxy $(C_1-C_6)$ alkylcarbonyloxy $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylcarbonyl,  $(C_1-C_6)$ alkylcarbonyl $(C_1-C_6)$ alkyl, carboxy,  $(C_1-C_6)$ alkoxycarbonyl,  $(C_8-C_{10})$ aryl $(C_1-C_6)$ alkoxycarbonyl,  $(C_8-C_{10})$ aryl $(C_1-C_6)$ alkylcarbonyl, aminocarbonyl,  $(C_1-C_6)$ alkylaminocarbonyl,  $((C_1-C_6)$ alkyl) $_2$ aminocarbonyl,  $(C_8-C_{10})$ arylaminocarbonyl,  $(C_8-C_{10})$ aryl $(C_1-C_6)$ alkylaminocarbonyl, carboxy $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxycarbonyl $(C_1-C_6)$ alkyl,  $(C_8-C_{10})$ aryl $(C_1-C_6)$ alkoxycarbonyl $(C_1-C_6)$ alkyl, aminocarbonyl $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylaminocarbonyl $(C_1-C_6)$ alkyl,  $((C_1-C_6)$ alkyl) $_2$ aminocarbonyl $(C_1-C_6)$ alkyl,  $(C_8-C_{10})$ arylaminocarbonyl $(C_1-C_6)$ alkyl,  $(C_8-C_{10})$ aryl $(C_1-C_6)$ alkylaminocarbonyl $(C_1-C_6)$ alkyl,  $(C_8-C_{10})$ arylsulfonyl,  $(C_2-C_9)$ heterocycloalkyl,  $(C_2-C_9)$ heteroaryl,  $(C_2-C_9)$ heterocycloalkyl $(C_1-$

C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl or R<sup>14</sup>R<sup>15</sup>N(C<sub>1</sub>-C<sub>6</sub>)alkyl wherein R<sup>14</sup> and R<sup>15</sup> are each independently (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl;

R<sup>4</sup> is (R<sup>5</sup>)<sub>f</sub>(R<sup>6</sup>)<sub>g</sub>(C<sub>6</sub>-C<sub>10</sub>)aryl, (R<sup>5</sup>)<sub>f</sub>(R<sup>6</sup>)<sub>g</sub>(C<sub>3</sub>-C<sub>10</sub>)cycloalkyl, (R<sup>5</sup>)<sub>f</sub>(R<sup>7</sup>)<sub>h</sub>(C<sub>2</sub>-C<sub>9</sub>)heteroaryl, or (R<sup>5</sup>)<sub>f</sub>(R<sup>7</sup>)<sub>h</sub>(C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl,

5 wherein f is 1, 2, 3 or 4;

g and h are each independently 0, 1, 2 or 3;

R<sup>5</sup> is one to three groups independently selected from (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylcarbonyl, (C<sub>2</sub>-C<sub>9</sub>)heteroarylcarbonyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylureido, amino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, acetylamino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, halo(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, acetylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, aminosulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminosulfonylamino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>2</sub>-C<sub>9</sub>)heteroaryloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, cyanoguanidino, (C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>cyanoguanidino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylcyanoguanidino, (C<sub>2</sub>-C<sub>9</sub>)heteroarylcyanoguanidino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino,

- amino(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, acetylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyano(C<sub>1</sub>-C<sub>6</sub>)alkylaminoalkyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, acetylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonyloxy(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyano(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyano(C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl,

- acetylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl,
- 5 C<sub>6</sub>)alkyl, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminosulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminosulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(cyanoguanidino)amino, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminosulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminosulfonyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylsulfonyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>2</sub>-C<sub>9</sub>)heteroarylaminosulfonyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylaminosulfonyl, R<sup>6</sup> is one to three groups independently selected from hydrogen, hydroxy, hydroxysulfonyl, halo, (C<sub>1</sub>-C<sub>6</sub>)alkyl, mercapto, mercapto(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylthio, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl, (C<sub>6</sub>-C<sub>10</sub>)arylsulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylthio(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>6</sub>-C<sub>10</sub>)aryloxy, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, trifluoro(C<sub>1</sub>-C<sub>6</sub>)alkyl, formyl, formyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, nitro, nitroso, cyano, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, trifluoro(C<sub>1</sub>-C<sub>6</sub>)alkoxy, amino(C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>3</sub>-C<sub>10</sub>)cycloalkyl,

- (C<sub>3</sub>-C<sub>10</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>3</sub>-C<sub>10</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>10</sub>)cycloalkylamino, (C<sub>3</sub>-C<sub>10</sub>)cycloalkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyano(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (hydroxy)(C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkylamino)(C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylthio(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino, (C<sub>6</sub>-C<sub>10</sub>)arylamino, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>2</sub>-C<sub>6</sub>)heterocycloalkylamino, (C<sub>2</sub>-C<sub>6</sub>)heteroarylamino, (C<sub>3</sub>-C<sub>10</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino, (C<sub>2</sub>-C<sub>6</sub>)alkenylcarbonylamino, (C<sub>3</sub>-C<sub>10</sub>)cycloalkylcarbonylamino, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonylamino, (C<sub>2</sub>-C<sub>6</sub>)heterocycloalkylcarbonylamino, halo(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, ((C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>10</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>6</sub>-C<sub>10</sub>)arylsulfonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)heterocycloalkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)heteroarylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl, (C<sub>6</sub>-C<sub>10</sub>)arylaminocarbonyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (amino(C<sub>1</sub>-C<sub>6</sub>)alkyl)aminocarbonyl, (hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, amidino, hydroxyamidino, guanidino, ureido, (C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>6</sub>-C<sub>10</sub>)arylureido, ((C<sub>6</sub>-C<sub>10</sub>)aryl)<sub>2</sub>ureido, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylureido,

halo(C<sub>1</sub>-C<sub>6</sub>)alkylureido, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)((C<sub>6</sub>-C<sub>10</sub>)aryl)ureido, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido, halo(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylureido, ureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (halo(C<sub>1</sub>-C<sub>6</sub>)alkyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)ureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl)ureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, glycinamido, (C<sub>1</sub>-C<sub>6</sub>)alkylglycinamido, aminocarbonylglycinamido, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylglycinamido, (aminocarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)glycinamido, ((C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)glycinamido, ((C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)glycinamido, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonylglycinamido, ((C<sub>6</sub>-C<sub>10</sub>)arylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)glycinamido, ((C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl)glycinamido, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)glycinamido, (C<sub>6</sub>-C<sub>10</sub>)arylaminocarbonylglycinamido, ((C<sub>6</sub>-C<sub>10</sub>)arylaminocarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)glycinamido, glycinamido(C<sub>1</sub>-C<sub>6</sub>)alkyl, alaninamido, (C<sub>1</sub>-C<sub>6</sub>)alkylalaninamido, alaninamido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl and (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sup>7</sup> is one to three groups independently selected from hydrogen, hydroxy, halo, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl, (C<sub>6</sub>-C<sub>10</sub>)arylsulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkoxy, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, fonyl, nitro, cyano, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino, (C<sub>6</sub>-C<sub>10</sub>)arylamino, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino, (C<sub>2</sub>-C<sub>6</sub>)alkenylcarbonylamino, cycloalkylcarbonylamino, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonylamino, halo(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, ((C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, aminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl, (C<sub>6</sub>-C<sub>10</sub>)arylaminocarbonyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, guanidino, ureido, (C<sub>1</sub>-C<sub>6</sub>)alkylureido, ureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, and glycinamido;


R<sup>8</sup> and R<sup>10</sup> are each independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl,



aminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl and (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl; and

- R<sup>11</sup> and R<sup>12</sup> are each independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>3</sub>-C<sub>8</sub>)cycloalkylcarbonylamino, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, ((C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroarylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino, halo(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonylamino, aminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, carboxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl and (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl;


wherein:



wherein the compound of formula X is converted to the corresponding compound of formula I by reacting X with a compound of the formula,  $H-(Z)_e-R^4$  wherein e is 1 and Z is oxygen, in the presence of potassium carbonate, potassium iodide and an aprotic solvent, such as butanone and the reaction is heated to reflux for a time period between about 4 hours to about 8 hours.


5

2. the method of preparation of claim 1, wherein:



wherein the compound of formula XXX is converted to the corresponding compound of formula I by reacting XXX with a compound of the formula, A-(X)<sub>c</sub>-(Y)<sub>d</sub>-(Z)<sub>e</sub>-R<sup>4</sup>, wherein A is chloro or bromo, in the presence of a base, such as triethylamine, and a polar aprotic solvent, such as methylene chloride and the reaction is stirred at a temperature between about -10°C to about 10°C, for a time period between about 15 minutes to about 90 minutes.


3. The method of preparation of claim 1, wherein:



wherein the compound of formula **XI** is converted to the corresponding compound of formula **II**, by reacting **XI** with an amine, in the presence of 4-dimethylaminopyridine, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide and a polar aprotic solvent, such as methylene chloride and the resulting reaction mixture is stirred overnight at room temperature.

4. The method of preparation of claim 1, wherein:


-76-



- wherein, for urea formation, the compound of formula **XIV** is converted to the corresponding compound of formula **V** by first reacting **XIV** with 4-nitrophenyl chloroformate in the presence of a base, such as pyridine, and a polar aprotic solvent, such as methylene chloride, followed by reacting the intermediate so formed with an amine and the reaction mixture, so formed, is allowed to stir overnight at room temperature; for sulfonamide formation, the compound of formula **XIV** is reacted with a sulfonyl chloride compound of the formula,  $\text{R}^{18}\text{-Cl}$ , in the presence of a base, such as triethylamine, and a polar aprotic solvent, such as methylene chloride and the reaction is stirred overnight at ambient temperature; for cyanoguanidine formation, the compound of formula **XIV** is first treated with sodium hydride in an aprotic solvent, such as tetrahydrofuran, followed by reacting, the intermediate so formed with dimethyl-N-cyanodithio iminocarbonate and the reaction mixture so formed is heated to reflux overnight and the N-cyano-S-methyl-isothioureia intermediate is then reacted with an amine in the presence of a polar protic solvent, such as methanol; for amide formation, the compound of formula **XIV** is reacted with an acid, such as 3-tert-butoxycarbonylaminopropionic acid in the presence of N-methylmorpholine, O-benzotriazole-1-yl-N,N,N', N'-tetramethyluronium hexafluorophosphate and a polar aprotic solvent, such as methylene chloride.

5. The method of preparation of claim 1, wherein:


-77-



- wherein the compound of formula **XVI** is converted to the corresponding compound of formula **VII** by reacting **XVI** with an amine of the formula,  $R^{16} R^{17}N$ , wherein  $R^{16}$  and  $R^{17}$  are each independently hydrogen, a nitrogen containing (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl or (C<sub>2</sub>-C<sub>9</sub>)heteroaryl group, or (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted by hydroxy, aminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>carbonyl, carboxy, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino, aminosulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminosulfonyl, (C<sub>6</sub>-C<sub>10</sub>)alkoxy, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, cyano, ureido, (C<sub>1</sub>-C<sub>6</sub>)alkylureido, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido, cyanoguanidino, (C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino and ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>cyanoguanidino, in the presence of a 10:1 ratio solution of dichloroethane/acetic acid and the reaction mixture is stirred, at room temperature, for a time period between about 30 minutes to about 2 hours and a reducing agent, such as sodium cyanoborohydride is then added to the mixture and the reaction is allowed to stir overnight at room temperature; when  $R^{16}$  and/or  $R^{17}$  is/are hydrogen, the compound of formula **VII** may further be reacted according to the procedure in claim, to provide ureas, sulfonamides, cyanoguanidinos, or amides.



6. The method of preparation of claim 1, wherein:

-78-



wherein the compound of formula **X** is converted to the corresponding compound of formula **XXXX** according to the procedure in claim 1 or the compound of formula **X** is converted to the corresponding compound of formula **XXXIX** according to the procedure in claim 1.

- 5                      7.        The method of preparation of claim 1, wherein:



wherein the amide compound of formula **XXXII** is converted to the corresponding compound of formula **XXXI** by reacting **XXXII** with a (C<sub>2</sub>-C<sub>9</sub>)heteroaryl lithium reagent in the presence of a polar aprotic solvent at a temperature between about -100°C to room temperature and the

5 resulting reaction mixture is stirred for a time period between about 1 hour to about 24 hours at a temperature between about -78°C to about 50°C.



(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
4 October 2001 (04.10.2001)

PCT

(10) International Publication Number  
WO 01/072728 A3

(51) International Patent Classification<sup>7</sup>: C07D 295/185,  
A61K 31/495, A61P 29/00, C07D 295/18, 213/40, 213/38,  
295/20, 307/20, 307/52, 207/16, 233/54, 307/14, 307/24,  
307/68

Paul [US/US]; Pfizer Global Research & Development,  
Eastern Point Road, Groton, CT 06340 (US).

(21) International Application Number: PCT/IB01/00375

(74) Agents: LUMB, J., Trevor et al.; Pfizer Inc., 235 East  
41nd Street, New York, NY 10017 (US).

(22) International Filing Date: 14 March 2001 (14.03.2001)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,  
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,  
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,  
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,  
TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/193,789 31 March 2000 (31.03.2000) US

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): PFIZER  
PRODUCTS INC. [US/US]; Eastern Point Road, Groton,  
CT 06340 (US).

(72) Inventors; and

**Published:**

- with international search report
- before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments


(75) Inventors/Applicants (*for US only*): BLUMBERG,  
Laura, Cook [US/US]; Pfizer Global Research & De-  
velopment, Eastern Point Road, Groton, CT 06340 (US).  
BROWN, Matthew, Frank [US/US]; Pfizer Global Re-  
search & Development, Eastern Point Road, Groton, CT  
06340 (US). MCGLYNN, Molly, Ann [US/US]; Pfizer  
Global Research & Development, Eastern Point Road,  
Groton, CT 06340 (US). POSS, Christopher, Stanley  
[US/US]; Pfizer Global Research & Development, Eastern  
Point Road, Groton, CT 06340 (US). GLADUE, Ronald,

(88) Date of publication of the international search report:  
18 July 2002

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*

WO 01/072728 A3

(54) Title: NOVEL PIPERAZINE DERIVATIVES



(57) Abstract: A compound of  
formula (I) or the pharmaceutically  
acceptable salt thereof; wherein a,  
b, c, d, e, j, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, and R<sup>4</sup> ar  
e as defined above useful to treat  
inflammation and other immune  
disorders.

# IN RNATIONAL SEARCH REPORT

International Application No

PCT/IB 01/00375

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D295/185 A61K31/495 A61P29/00 C07D295/18 C07D213/40  
 C07D213/38 C07D295/20 C07D307/20 C07D307/52 C07D207/16  
 C07D233/54 C07D307/14 C07D307/24 C07D307/68

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, EP0-Internal

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 56771 A (SCHERING AG) 17 December 1998 (1998-12-17) page 75 -page 98; claims -----	1-7

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*G\* document member of the same patent family

Date of the actual completion of the international search

10 May 2002

Date of mailing of the international search report

16/05/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Sánchez García, J.M.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 01/00375

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9856771	A	17-12-1998	US 6207665 B1 27-03-2001
			AU 735462 B2 12-07-2001
			AU 8625898 A 30-12-1998
			EE 9900565 A 15-06-2000
			WO 9856771 A2 17-12-1998
			EP 0988292 A2 29-03-2000
			HU 0003929 A2 28-05-2001
			JP 2002503239 T 29-01-2002
			NO 996068 A 11-02-2000
			PL 337758 A1 11-09-2000
			SK 171399 A3 12-09-2000
			TR 9903034 T2 21-06-2000

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
4 October 2001 (04.10.2001)

PCT

(10) International Publication Number  
**WO 01/072728 A3**(51) International Patent Classification<sup>7</sup>: C07D 295/185, A61K 31/495, A61P 29/00, C07D 295/18, 213/40, 213/38, 295/20, 307/20, 307/52, 207/16, 233/54, 307/14, 307/24, 307/68

(21) International Application Number: PCT/IB01/00375

(22) International Filing Date: 14 March 2001 (14.03.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/193,789 31 March 2000 (31.03.2000) US

(74) Agents: LUMB, J., Trevor et al.; Pfizer Inc., 235 East 41nd Street, New York, NY 10017 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).(71) Applicant (*for all designated States except US*): PFIZER PRODUCTS INC. [US/US]; Eastern Point Road, Groton, CT 06340 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): BLUMBERG, Laura, Cook [US/US]; Pfizer Global Research & Development, Eastern Point Road, Groton, CT 06340 (US). BROWN, Matthew, Frank [US/US]; Pfizer Global Research & Development, Eastern Point Road, Groton, CT 06340 (US). MCGLYNN, Molly, Ann [US/US]; Pfizer Global Research & Development, Eastern Point Road, Groton, CT 06340 (US). POSS, Christopher, Stanley [US/US]; Pfizer Global Research & Development, Eastern Point Road, Groton, CT 06340 (US). GLADUE, Ronald, Paul [US/US]; Pfizer Global Research & Development, Eastern Point Road, Groton, CT 06340 (US).

Published:

- with international search report
- with amended claims

(88) Date of publication of the international search report:  
18 July 2002

Date of publication of the amended claims: 9 January 2003

(15) Information about Correction:


Previous Correction:

see PCT Gazette No. 41/2002 of 10 October 2002, Section II

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/072728 A3

(54) Title: NOVEL PIPERAZINE DERIVATIVES




(1)

(57) Abstract: A compound of formula (I) or the pharmaceutically acceptable salt thereof; wherein a, b, c, d, e, j, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, and R<sup>4</sup> are as defined above useful to treat inflammation and other immune disorders.

## AMENDED CLAIMS

[received by the International Bureau on 8 July 2002 (08.07.02);  
original claims 1-7 repaced by amended claims 1-13 (12 pages)]

1. A compound of the formula



or the pharmaceutically acceptable salt thereof; wherein

- 5           a is 1, 2, 3, 4 or 5;  
          b is 0, 1, 2, 3 or 4;  
          c is 0 or 1;  
          d is 1, 2, 3, 4 or 5;  
          e is 0 or 1;
- 10          j is 1, 2, 3, or 4;  
          X is C(O), C(S) or CH<sub>2</sub>;  
          Y is CH<sub>2</sub>, or if e is 0, Y is CHR<sup>8</sup> wherein R<sup>8</sup> is hydrogen, (C<sub>6</sub>-C<sub>10</sub>)aryl or NR<sup>9</sup>R<sup>10</sup>;  
          Z is oxygen, NR<sup>9</sup> or CR<sup>11</sup>R<sup>12</sup>;
- each R<sup>1</sup> is independently selected from hydrogen, hydroxy, hydroxysulfonyl, halo,
- 15   (C<sub>1</sub>-C<sub>6</sub>)alkyl, mercapto, mercapto(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylthio, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylthio(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>6</sub>-C<sub>10</sub>)aryloxy, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, trifluoromethyl, formyl, formyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, nitro, nitroso, cyano, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, trifluoromethoxy, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkylamino, (C<sub>3</sub>-C<sub>7</sub>)cycloalkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>3</sub>-C<sub>7</sub>)cycloalkyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, ((C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyano(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>7</sub>)alkenyl, (C<sub>2</sub>-C<sub>7</sub>)alkynyl, (C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>2</sub>-C<sub>8</sub>)alkenyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylthio(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino, (C<sub>6</sub>-C<sub>10</sub>)arylamino, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, carboxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-

$C_{10}$ )arylcarbonyl,  $(C_6-C_{10})$ arylcarbonyl $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkylcarbonyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkylcarbonyl $(C_1-C_6)$ alkyl, carboxy $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxycarbonyl $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkoxycarbonyl $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxy $(C_1-C_6)$ alkylcarbonyloxy $(C_1-C_6)$ alkyl, aminocarbonyl,  $(C_1-C_6)$ alkylaminocarbonyl,  $((C_1-C_6)$ alkyl)<sub>2</sub>aminocarbonyl,  $(C_6-C_{10})$ arylaminocarbonyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkylaminocarbonyl, aminocarbonyl $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylaminocarbonyl $(C_1-C_6)$ alkyl,  $((C_1-C_6)$ alkyl)<sub>2</sub>aminocarbonyl $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ arylaminocarbonyl $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylaminocarbonyl $(C_1-C_6)$ alkyl, amidino, guanidino, ureido,  $(C_1-C_6)$ alkylureido,  $((C_1-C_6)$ alkyl)<sub>2</sub>ureido, ureido $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylureido $(C_1-C_6)$ alkyl,  $((C_1-C_6)$ alkyl)<sub>2</sub>ureido $(C_1-C_6)$ alkyl,  $(C_2-C_9)$ heterocycloalkyl,  $(C_2-C_9)$ heteroaryl,  $(C_2-C_9)$ heterocycloalkyl $(C_1-C_6)$ alkyl and  $(C_2-C_9)$ heteroaryl $(C_1-C_6)$ alkyl;

each  $R^2$  and  $R^3$  are independently selected from oxo, halo,  $(C_1-C_6)$ alkyl,  $(C_3-C_8)$ cycloalkyl,  $(C_3-C_8)$ cycloalkyl $(C_1-C_6)$ alkyl,  $(C_3-C_8)$ cycloalkylamino $(C_1-C_6)$ alkyl,  $(C_3-C_8)$ cycloalkyl $(C_1-C_6)$ alkylamino $(C_1-C_6)$ alkyl, halo $(C_1-C_6)$ alkyl,  $(C_2-C_6)$ alkenyl,  $(C_2-C_6)$ alkynyl,  $(C_6-C_{10})$ aryl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl $(C_2-C_6)$ alkenyl, H-C(O)-, H-C(O)- $(C_1-C_6)$ alkyl, hydroxy $(C_1-C_6)$ alkyl, hydroxy $(C_2-C_6)$ alkenyl, hydroxy $(C_2-C_6)$ alkynyl, hydroxy $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkyl, hydroxy $(C_3-C_8)$ cycloalkyl $(C_1-C_6)$ alkyl, thio $(C_1-C_6)$ alkyl, cyano $(C_1-C_6)$ alkyl, halo $(C_1-C_6)$ alkylcarbonylamino $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxy $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxy $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryloxy $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkoxy $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylthio $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylsulfinyl $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylsulfonyl $(C_1-C_6)$ alkyl, hydroxy $(C_1-C_6)$ alkylthio $(C_1-C_6)$ alkyl, amino $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylamino $(C_1-C_6)$ alkyl,  $((C_1-C_6)$ alkyl)<sub>2</sub>amino $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ arylamino $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkylamino $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylcarbonylamino $(C_1-C_6)$ alkyl, azido $(C_1-C_6)$ alkyl, aminocarbonylamino $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylaminocarbonylamino $(C_1-C_6)$ alkyl,  $((C_1-C_6)$ alkyl)<sub>2</sub>aminocarbonylamino $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxycarbonyl $(C_1-C_6)$ alkylaminocarbonylamino $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxycarbonylamino $(C_1-C_6)$ alkyl, hydroxy $(C_1-C_6)$ alkylamino $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryloxy $(C_1-C_6)$ alkylcarbonyloxy $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxy $(C_1-C_6)$ alkylcarbonyloxy $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkoxy $(C_1-C_6)$ alkylcarbonyloxy $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylcarbonyl,  $(C_1-C_6)$ alkylcarbonyl $(C_1-C_6)$ alkyl, carboxy,  $(C_1-C_6)$ alkoxycarbonyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkoxycarbonyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkylcarbonyl, aminocarbonyl,  $(C_1-C_6)$ alkylaminocarbonyl,  $((C_1-C_6)$ alkyl)<sub>2</sub>aminocarbonyl,  $(C_6-C_{10})$ arylaminocarbonyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkylaminocarbonyl, carboxy $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxycarbonyl $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkoxycarbonyl $(C_1-C_6)$ alkyl, aminocarbonyl $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylaminocarbonyl $(C_1-C_6)$ alkyl,  $((C_1-C_6)$ alkyl)<sub>2</sub>aminocarbonyl $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ arylaminocarbonyl $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkylaminocarbonyl $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ arylsulfonyl,  $(C_2-C_9)$ heterocycloalkyl,  $(C_2-C_9)$ heteroaryl,  $(C_2-C_9)$ heterocycloalkyl $(C_1-C_6)$ alkyl and  $(C_2-C_9)$ heteroaryl $(C_1-C_6)$ alkyl.

C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl or R<sup>14</sup>R<sup>15</sup>N(C<sub>1</sub>-C<sub>6</sub>)alkyl wherein R<sup>14</sup> and R<sup>15</sup> are each independently (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl;

R<sup>4</sup> is (R<sup>5</sup>)<sub>f</sub>(R<sup>6</sup>)<sub>g</sub>(C<sub>6</sub>-C<sub>10</sub>)aryl, (R<sup>5</sup>)<sub>f</sub>(R<sup>6</sup>)<sub>g</sub>(C<sub>3</sub>-C<sub>10</sub>)cycloalkyl, (R<sup>5</sup>)<sub>f</sub>(R<sup>7</sup>)<sub>h</sub>(C<sub>2</sub>-C<sub>9</sub>)heteroaryl, or (R<sup>5</sup>)<sub>f</sub>(R<sup>7</sup>)<sub>h</sub>(C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl,

5 wherein f is 1, 2, 3 or 4;

g and h are each independently 0, 1, 2 or 3;

R<sup>5</sup> is one to three groups independently selected from (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylcarbonyl, (C<sub>2</sub>-C<sub>9</sub>)heteroarylcarbonyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylureido, amino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, acetylmino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, halo(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, acetylmino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, aminosulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminosulfonylamino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>2</sub>-C<sub>9</sub>)heteroaryloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, cyanoguanidino, (C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>cyanoguanidino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylcyanoguanidino, (C<sub>2</sub>-C<sub>9</sub>)heteroarylcyanoguanidino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino,

- amino(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, acetylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyano(C<sub>1</sub>-C<sub>6</sub>)alkylaminoalkyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, acetylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonyloxy(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyano(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyano(C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl,



- acetylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl,
- 5 C<sub>6</sub>)alkyl, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminosulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminosulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(cyanoguanidino)amino, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminosulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminosulfonyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylsulfonyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>2</sub>-C<sub>9</sub>)heteroarylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylaminosulfonyl, R<sup>8</sup> is one to three groups independently selected from hydrogen, hydroxy, hydroxysulfonyl, halo, (C<sub>1</sub>-C<sub>6</sub>)alkyl, mercapto, mercapto(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylthio, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl, (C<sub>6</sub>-C<sub>10</sub>)arylulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylthio(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>6</sub>-C<sub>10</sub>)aryloxy, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, trifluoro(C<sub>1</sub>-C<sub>6</sub>)alkyl, formyl, formyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, nitro, nitroso, cyano, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, trifluoro(C<sub>1</sub>-C<sub>6</sub>)alkoxy, amino(C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>3</sub>-C<sub>10</sub>)cycloalkyl,

- (C<sub>3</sub>-C<sub>10</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>3</sub>-C<sub>10</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>10</sub>)cycloalkylamino, (C<sub>3</sub>-C<sub>10</sub>)cycloalkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyano(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (hydroxy)(C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkylamino)(C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylthio(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino, (C<sub>6</sub>-C<sub>10</sub>)arylamino, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylamino, (C<sub>2</sub>-C<sub>9</sub>)heteroarylamino, (C<sub>3</sub>-C<sub>10</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino, (C<sub>2</sub>-C<sub>6</sub>)alkenylcarbonylamino, (C<sub>3</sub>-C<sub>10</sub>)cycloalkylcarbonylamino, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonylamino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylcarbonylamino, halo(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, ((C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>10</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroarylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl, (C<sub>6</sub>-C<sub>10</sub>)arylaminocarbonyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (amino(C<sub>1</sub>-C<sub>6</sub>)alkyl)aminocarbonyl, (hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, amidino, hydroxyamidino, guanidino, ureido, (C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>6</sub>-C<sub>10</sub>)arylureido, ((C<sub>6</sub>-C<sub>10</sub>)aryl)<sub>2</sub>ureido, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylureido,

halo(C<sub>1</sub>-C<sub>6</sub>)alkylureido, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)((C<sub>6</sub>-C<sub>10</sub>)aryl)ureido, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido, halo(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylureido, ureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (halo(C<sub>1</sub>-C<sub>6</sub>)alkyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)ureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl)ureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, glycinamido, (C<sub>1</sub>-C<sub>6</sub>)alkylglycinamido, aminocarbonylglycinamido, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylglycinamido, (aminocarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)glycinamido, ((C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)glycinamido, ((C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)glycinamido, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonylglycinamido, ((C<sub>6</sub>-C<sub>10</sub>)arylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)glycinamido, ((C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl)glycinamido, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)glycinamido, (C<sub>6</sub>-C<sub>10</sub>)arylaminocarbonylglycinamido, ((C<sub>6</sub>-C<sub>10</sub>)arylaminocarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)glycinamido, glycinamido(C<sub>1</sub>-C<sub>6</sub>)alkyl, alaninamido, (C<sub>1</sub>-C<sub>6</sub>)alkylalaninamido, alaninamido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl and (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sup>7</sup> is one to three groups independently selected from hydrogen, hydroxy, halo, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl, (C<sub>6</sub>-C<sub>10</sub>)arylsulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkoxy, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, formyl, nitro, cyano, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino, (C<sub>6</sub>-C<sub>10</sub>)arylamino, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino, (C<sub>2</sub>-C<sub>6</sub>)alkenylcarbonylamino, cycloalkylcarbonylamino, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonylamino, halo(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, ((C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, aminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl, (C<sub>6</sub>-C<sub>10</sub>)arylaminocarbonyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, guanidino, ureido, (C<sub>1</sub>-C<sub>6</sub>)alkylureido, ureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, and glycinamido;

R<sup>9</sup> and R<sup>10</sup> are each independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl,

aminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl and (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl; and

R<sup>11</sup> and R<sup>12</sup> are each independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>3</sub>-C<sub>8</sub>)cycloalkylcarbonylamino, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>6</sub>-C<sub>10</sub>)arylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, ((C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroarylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino, halo(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonylamino, aminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, carboxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl and (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl.

2. A compound according to claim 1, wherein R<sup>1</sup> is hydrogen, halo, cyano, nitro, trifluoromethyl, trifluoromethoxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy or (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyloxy.

3. A compound according to claim 1, wherein R<sup>2</sup> and R<sup>3</sup> are each independently selected from (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl.

4. A compound according to claim 1, wherein c is 1; X is C(O) or CH<sub>2</sub>; d is 2; Y is ethylene; and e is 0.

5. A compound according to claim 1, wherein c is 1; X is C(O) or CH<sub>2</sub>; d is 1 or 2; Y is CH<sub>2</sub> or ethylene; e is 1; and Z is oxygen or NR<sup>9</sup> wherein R<sup>9</sup> is hydrogen or (C<sub>1</sub>-C<sub>6</sub>)alkyl.

6. A compound according to claim 1, wherein c is 1; X is C(O) or CH<sub>2</sub>; d is 1; Y is CHR<sup>8</sup> wherein R<sup>8</sup> is NR<sup>9</sup>R<sup>10</sup>; R<sup>9</sup> and R<sup>10</sup> are each independently hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl; e is 1; and Z is selected from the group consisting of oxygen; CR<sup>11</sup>R<sup>12</sup> wherein R<sup>11</sup> and R<sup>12</sup> are hydrogen, and NR<sup>9</sup> wherein R<sup>9</sup> is hydrogen or (C<sub>1</sub>-C<sub>6</sub>)alkyl.

7. A compound according to claim 1, wherein R<sup>4</sup> is (R<sup>5</sup>)<sub>f</sub>(R<sup>6</sup>)<sub>g</sub>(C<sub>6</sub>-C<sub>10</sub>)aryl or (R<sup>5</sup>)<sub>f</sub>(R<sup>7</sup>)<sub>h</sub>(C<sub>2</sub>-C<sub>9</sub>)heteroaryl wherein f, g and h are independently 1 or 2.

8. A compound according to claim 1, wherein R<sup>5</sup> is selected from the group consisting of (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylcarbonyl, (C<sub>2</sub>-C<sub>9</sub>)heteroarylcarbonyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, aminosulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylureido, acetilamino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylureido, amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, acetilamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylcyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, aminosulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminosulfonylamino, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>2</sub>-C<sub>9</sub>)heteroaryloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, amino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl amino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyloxycarbonyl amino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-

- C<sub>6</sub>alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, acetylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (acetyl)((C<sub>1</sub>-C<sub>6</sub>)alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl,
- 5 C<sub>6</sub>alkyl, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>(cyanoguanidino)(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl,
- 10 aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heteroarylloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminosulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminosulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylsulfonyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>2</sub>-C<sub>9</sub>)heteroarylaminosulfonyl, ureido(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylureido(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>ureido(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkyloxycarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, aminocarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, cyanoguanidino(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>2</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, (C<sub>2</sub>-C<sub>9</sub>)heterocycloalkylaminosulfonyl halo(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylureido, halo(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylaminocarbonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, aminosulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>aminosulfonyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl, and (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkylaminosulfonyl.

9. A compound according to claim 1, wherein R<sup>6</sup> and R<sup>7</sup> are each independently  
 30 halo, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, trifluoromethyl, trifluoromethoxy, hydroxy, aminocarbonyl, cyano, ureido, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylamino or glycinamino.

10. A pharmaceutical composition for treating or preventing a disorder or condition  
 35 selected from autoimmune diseases, rheumatoid arthritis, type I diabetes (recent onset), lupus, inflammatory bowel disease, optic neuritis, psoriasis, multiple sclerosis, polymyalgia rheumatica, uveitis, and vasculitis, acute and chronic inflammatory conditions osteoarthritis, adult Respiratory Distress Syndrome, Respiratory Distress Syndrome of infancy, ischemia

reperfusion injury, glomerulonephritis, and chronic obstructive pulmonary disease (COPD) allergic conditions, asthma and atopic dermatitis, inflammation associated with infection, viral inflammation, influenza, hepatitis and Guillian-Barre, chronic bronchitis, chronic or acute tissue, cell, and solid organ transplant rejection, xeno-transplantation, atherosclerosis, restenosis, HIV infectivity (co-receptor usage), and granulomatous diseases, sarcoidosis, leprosy and tuberculosis, and sequelae associated with cancers, multiple myeloma; limiting the production of cytokines and/or TNF at inflammatory sites, as a consequence of decreasing cell infiltration; for treating diseases and/or congestive heart failure, linked to TNF and IL-1 and for treating pulmonary emphysema or dyspnea associated therewith, emphysema; HIV-1, HIV-2, HIV-3; cytomegalovirus (CMV), adenoviruses; Herpes viruses (*Herpes zoster* and *Herpes simplex*), for treating sequelae associated with infection where such infection induces production of detrimental inflammatory cytokines and/or TNF, fungal meningitis, joint tissue damage, hyperplasia, pannus formation and bone resorption, psoriatic arthritis, hepatic failure, bacterial meningitis, Kawasaki syndrome, myocardial infarction, acute liver failure, Lyme disease, septic shock, cancer, trauma, and malaria, in a mammal, comprising an amount of a compound according to claim 1, or a pharmaceutically acceptable salt or pro-drug thereof, that is effective in treating or preventing such disorder or condition and a pharmaceutically acceptable carrier.

11. A pharmaceutical composition for treating or preventing a disorder or condition that can be treated or prevented by inhibiting chemokine binding to the receptor CCR1 in a mammal, comprising an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, effective in treating or preventing such disorder or condition and a pharmaceutically acceptable carrier.

12. A method for treating or preventing a disorder or condition selected from autoimmune diseases, rheumatoid arthritis, type I diabetes (recent onset), lupus, inflammatory bowel disease, optic neuritis, psoriasis, multiple sclerosis, polymyalgia rheumatica, uveitis, and vasculitis, acute and chronic inflammatory conditions osteoarthritis, adult Respiratory Distress Syndrome, Respiratory Distress Syndrome of infancy, ischemia reperfusion injury, glomerulonephritis, and chronic obstructive pulmonary disease (COPD) allergic conditions, asthma and atopic dermatitis, inflammation associated with infection, viral inflammation, influenza, hepatitis and Guillian-Barre, chronic bronchitis, chronic or acute tissue, cell, and solid organ transplant rejection, xeno-transplantation, atherosclerosis, restenosis, HIV infectivity (co-receptor usage), and granulomatous diseases, sarcoidosis, leprosy and tuberculosis, and sequelae associated with cancers, multiple myeloma; limiting the production of cytokines and/or TNF at inflammatory sites, as a consequence of decreasing cell infiltration; for treating diseases and/or congestive heart failure, linked to TNF and IL-1 and for treating pulmonary emphysema or dyspnea associated therewith,

- emphysema; HIV-1, HIV-2, HIV-3; cytomegalovirus (CMV), adenoviruses, Herpes viruses (*Herpes zoster* and *Herpes simplex*), for treating sequelae associated with infection where such infection induces production of detrimental inflammatory cytokines and/or TNF, fungal meningitis, joint tissue damage, hyperplasia, pannus formation and bone resorption, psoriatic
- 5 arthritis, hepatic failure, bacterial meningitis, Kawasaki syndrome, myocardial infarction, acute liver failure, lyme disease, septic shock, cancer, trauma, and malaria, in a mammal, comprising administering to a mammal in need of such treatment or prevention an amount of a compound according to claim 1, or a pharmaceutically acceptable salt or pro-drug thereof, that is effective in treating or preventing such disorder or condition.
- 10 13. A method for treating or preventing a disorder or condition that can be treated or prevented by antagonizing the CCR1 receptor in a mammal, comprising administering to a mammal in need of such treatment or prevention an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder or condition.